

Bibliography of the Sources Examined in preparation of the Opinion

- Altenburger R, Boedeker W, Faust M, Grimme LH (1996). Regulations for combined effects of pollutants: consequences from risk assessment in aquatic toxicology. *Food Chem Toxicol* 34, 1155-1157
- Altenburger R, Walter H, Grote M (2004). What Contributes to the Combined Effect of a Complex Mixture? *Environ Sci Technol.* 38, 6353-6362
- Ankley GT, Bennett RS, Erickson RJ et al. (2010) Adverse outcome pathways: a conceptual framework to support ecotoxicology research and risk assessment. *Environ Toxicol Chem.* 2010 Mar;29(3):730-41
- Ankley GT., Bennett RS., Erickson RJ;, Hoff DJ;, Hornung MW., Johnson RD., Mount DR., Nichols JW., Russom CL., Schmieder PK., Serrano JA., Tietge JE., Villeneuve DL. (2010). Adverse outcome pathways: a conceptual framework to support ecotoxicology research and risk assessment. *Environmental Toxicology and Chemistry*, Vol. 29, No. 3, pp. 730–741, 2010
- Arts GHP, Buijse-Bogdan LL, Belgers JDM, van Rhenen-Kersten CH, van Wijngaarden, RPA, Roessink I, Maund SJ, van den Brink PJ, Brock TCM. (2006) Ecological Impact in Ditch Mesocosms of Simulated Spray Drift from a Crop Protection Program for Potatoes. *Integrated Environmental Assessment and Management* — Volume 2, Number 2—pp. 105–125
- ATSDR (2004). Guidance Manual for the Assessment of Joint Toxic Action of Chemical Mixtures. US Agency for Toxic Substances and Disease Registry. Division of Toxicology. May 2004
- Balk L., Hägerroth P-A., Åkerman G., Hanson M., Tjärnlund U., Hansson T., Hallgrimsson GT., Zebühr Y., Broman D., Mörner T., Sundberg H. (2009). Wild birds of declining European species are dying from a thiamine deficiency syndrome. *PNAS* July 21, 2009 vol. 106 no. 29 12001–12006. www.pnas.org/cgi/doi/10.1073/pnas.0902903106
- Balk L., Hägerroth P-A., Åkerman G., Hanson M., Tjärnlund U., Hansson T., Hallgrimsson GT., Zebühr Y., Broman D., Mörner T., Sundberg H. (2009). Supporting Information. Wild birds of declining European species are dying from a thiamine deficiency syndrome.
- BEAM (2003) Bridging Effect Assessment of Mixtures to Ecosystem Situations and Regulation – BEAM – Final Summary Report. Section 5 and 6; deliverable 17. July 2003. Contract No.: EVK1-CT-1999-00012. Coordinator: Prof. Dr. L. H. Grimme, Dr. T. Backhaus, University of Bremen
- BEAM Risk Assessment of Chemical Mixtures in the Aquatic Environment BEAM Workpackage 8 (Deliverables 13/14). Report. Dr. Thomas Backhaus, Dr. Michael Faust, Martin Scholze, Marion Junghans, Prof. Dr. L. H. Grimme, University of Bremen, Germany. Prof. Marco Vighi, University of Milano Bicocca, Italy. Dr. Rolf Altenburger, Centre for Environmental Research, Leipzig/Halle, Germany
- BEAM. Risk Assessment of Chemical Mixtures in the Aquatic Environment. BEAM Workpackage 8 (Deliverables 13/14). Report. Backhaus T., Faust M., Scholze M., Junghans M., Grimme LH., Vighi M., Altenburger T. 2003
- Benson R. (2009). Hazard to the developing male reproductive system from cumulative exposure to phthalate esters—dibutyl phthalate, diisobutyl phthalate, butylbenzyl phthalate, diethylhexyl phthalate, dipentyl phthalate, and diisononyl phthalate. *Regulatory Toxicology and Pharmacology* 53 (2009) 90–101
- Bliss CI (1939). The toxicity of poisons applied jointly. *Ann Appl Biol.* 26(3), 585-615
- Boobis AR, Budinsky R, Collie S, Crofton K, Embry M, Felter S, Hertzberg R, Kopp D, Mihlan G, Mumtaz M, Price P, Solomon K, Teuschler L, Yang R, Zaleski R (2011). Critical

- analysis of literature on low-dose synergy for use in screening chemical mixtures for risk assessment. *Crit Rev Toxicol* 2011; 41(5):369-83
- Boobis AR, Ossendorp BC, Banasiak U, Hamey PY, Sebestyen I, Moretto A (2008). Cumulative risk assessment of pesticide residues in food. *Toxicol Lett.* 180, 137-150
- Cao Z., Shafer TJ., Crofton KM., Gennings C., Murray TF. (2011) Additivity of Pyrethroid Actions on Sodium Influx in Cerebrocortical Neurons in Primary Culture. *Environ Health Perspect* 119:1239-1246 (2011). <http://dx.doi.org/10.1289/ehp.1003394> [Online 10 June 2011]
- Carlson-Lynch H, Price PS, Swartout JC, Dourson ML, Keenan RE (1999). Application of Quantitative Information on the Uncertainty in the RfD to Noncarcinogenic Risk Assessments. *Human and Ecological Risk Assessment* 5(3), 527-546
- Carmichael N., Bausen M., Boobis AR., Cohen SM;, Embry M., Fruijtjer-Pölloth C., Greim H., Lewis R., Meek M.E. (Bette), Mellor H., Vickers C., Doe J. (2011) .Using mode of action information to improve regulatory decision-making: An ECETOC/ILSI RF/HESI workshop overview. *Critical Reviews in Toxicology*, 2011; 41(3): 175-186
- CEFIC. Letter on proposed decision tree for the risk assessment of mixtures. 9 September 2011
- Charles GD, Gennings C, Tornesi B, Kan HL, Zacharewski TR, Bhaskar Gollapudi B, Carney EW (2007). Analysis of the interaction of phytoestrogens and synthetic chemicals: an in vitro/in vivo comparison. *Toxicol Appl Pharmacol.* 218(3), 280-288
- Charles GD, Gennings C, Zacharewski TR, Gollapudi BB, Carney EW (2002). An approach for assessing estrogen receptor-mediated interactions in mixtures of three chemicals: A pilot study. *Toxicol Sci.* 68(2), 349-360
- Chen CL, Hsu LI, Chiou HY, Hsueh YM, Chen SY, Wu MM et al. (2004). Ingested arsenic, cigarette smoking, and lung cancer risk: A follow-up study in arseniasis-endemic areas in Taiwan. *JAMA* 292, 2984-2990
- Christiansen S, Scholze M, Dalgaard M, Vinggaard AM, Axelstad M, Kortenkamp A, Hass U (2009). Synergistic Disruption of External Male Sex Organ Development by a Mixture of Four Antiandrogens. *Environ Health Perspect.* 117(12), 1839-1846
- Christiansen S., Scholze M., Dalgaard M., Vinggaard AM., Axelstad M., Kortenkamp A., Hass U. (2009). Synergistic Disruption of External Male Sex Organ Development by a Mixture of Four Antiandrogens. *Environ Health Perspect* 117:1839-1846 (2009)
- Collins FS., Gray GM., Bucher JR. (2008). Transforming Environmental Health Protection. 15 FEBRUARY 2008 VOL 319 SCIENCE PP. 906-907. www.sciencemag.org
- COT (2002). UK Committee on Toxicity of Chemicals in Food, Consumer Products and the Environment. *Risk Assessment of Mixtures of Pesticides and Similar Substances*
- Crofton K, Craft ES, Hedge JM, Gennings C, Simmons JE, Carchman RA, Carter WH Jr, de Vito JM (2005). Thyroid-hormone-disrupting chemicals: evidence for dose-dependent additivity or synergism. *Environ Health Perspect.* 113(11), 1549-1554
- Cronin MTD, Jaworska JS, Walker JD, Comber MHI, Watts CD, Worth AP (2003a). Use of QSARs in International Decision-Making Frameworks to Predict Health Effects of Chemical Substances. *Environ Health Perspect.* 111, 1391-1401
- Cronin MTD, Walker JD, Jaworska JS, Comber MHI, Watts CD, Worth AP (2003b). Use of QSARs in International Decision-Making Frameworks to Predict Ecologic Effects and Environmental Fate of Chemical Substances. *Environ Health Perspect.* 111, 1376-1390
- CSTEE 1999 (Scientific Committee for Toxicity, Ecotoxicity and the Environment. CSTEE) Opinion on human and wildlife health effects of endocrine disrupting chemicals, with emphasis on wildlife and on ecotoxicology test methods. March 1999

- CVUA (2007). Chemisches und Veterinäruntersuchungsamt Stuttgart/Germany. Toxikologische Bewertung von Mehrfachrückständen in Obst und Gemüse. Report
- De Wolf W., Siebel-Sauer A., Lecloux A., Koch V., Holt M., Feijtel T., Comber M., Boeije G. (2005). Mode of action and aquatic exposure thresholds of no concern. *Environmental Toxicology and Chemistry*, Vol. 24, No. 2, pp. 479–485, 2005
- De Zwart D., Posthuma L. (2005). Complex mixture toxicity for single and multiple species: proposed methodologies. *Environmental Toxicology and Chemistry*, Vol. 24, No. 10, pp. 2665–2676, 2005
- Desai AM, Autenrieth RL, Dimitriou-Christidis P, McDonald TJ (2008). Biodegradation kinetics of select polycyclic aromatic hydrocarbon (PAH) mixtures by *Sphingomonas paucimoblis* EPA505. *Biodegradation*, 19(2), 223-233
- Dimitriou-Christidis P, Autenrieth RL (2007). Kinetics of biodegradation of binary and ternary mixtures of PAHs. *Biotechnol Bioeng*. 97(4), 788-800
- ECHA (2008) European Chemicals Agency. Guidance on information requirements and chemical safety assessment. Chapter R.6: QSARs and grouping of chemicals. May 2008
http://guidance.echa.europa.eu/docs/guidance_document/information_requirements_en.htm?time=1323956054 (accessed on 14 December 2011).
- EFSA (2006). European Food Safety Authority. Opinion of the Scientific Committee related to uncertainties in dietary exposure assessment. *EFSA Journal* 438, 1-54
- EFSA (2008). European Food Safety Authority. Opinion of the Scientific Panel on Plant Protection products and their Residues to evaluate the suitability of existing methodologies and, if appropriate, the identification of new approaches to assess cumulative and synergistic risks from pesticides to human health with a view to set MRLs for those pesticides in the frame of Regulation (EC) 396/2005. *The EFSA Journal* 704, 1-84
- EFSA (2009). European Food Safety Authority. Panel on Plant Protection Products and their Residues (PPR Panel) Scientific Opinion on a request from EFSA on risk assessment for a selected group of pesticides from the triazole group to test possible methodologies to assess cumulative effects from exposure through food from these pesticides on human health. *The EFSA Journal* 7(9), 1167
- EFSA (2009a). European Food Safety Authority. Guidance of the Scientific Committee on a request from EFSA on the use of the benchmark dose approach in risk assessment. *The EFSA Journal* 1150, 1-72
- Faust M, Altenburger R, Backhaus T, Blanck H, Boedeker W, Gramatica P, Hamer V, Scholze M, Vighi M, Grimme LH (2003). Joint algal toxicity of 16 dissimilarly acting chemicals is predictable by the concept of independent action. *Aquat Toxicol* 63(1), 43-63
- Faust M, Altenburger R, Boedeker W, Grimme LH (1994). Algal toxicity of binary combinations of pesticides. *Bull Environ Contam Toxicol* 53, 134-141
- Feron VJ and Groten JP (2002). Toxicological evaluation of chemical mixtures. *Food Chem Toxicol* 40(6), 825-839
- Finizio A, Villa S, Vighi M (2005). Predicting pesticide mixtures in surface waters from a given crop. *Agriculture Ecosystems and Environment* 111, 111-118
- Gomes J and Meek B (2009). Interactions between Occupational and Environmental Factors in Toxicology, Hazard Evaluation and Risk Assessment. Book chapter in: *General, Applied and Systems Toxicology*. John Wiley & Sons, Ltd.
- Grandjean P., Landrigan PJ. (2006). Developmental neurotoxicity of industrial chemicals. www.thelancet.com Published online November 8, 2006 DOI:10.1016/S0140-6736(06)69665-7

- Greim H and Snyder R (2008). Toxicology and risk assessment: a comprehensive introduction. Wiley-Interscience
- Gross M., Daginnus K., Deviller G., de Wolf W., Dungey S., Galli C., Gourmelon A., Jacobs M., Matthiessen P., Micheletti C., Nestmann E., Pavan M., Paya-Perez A., Ratte H-T., Safford B., Sokull-Kluttgen B., Stock F., Stolzenberg H-C., Wheeler J., Willuhn M., Worth A., Zaldivar Comenges JM., Crane M. (2009). Thresholds of Toxicological Concern for Endocrine Active Substances in the Aquatic Environment. Integrated Environmental Assessment and Management – Volume 6, Number 1—pp. 2–11. 2009
- Guillette LJ., Edwards TM., Moore BC. (2007) Alligators, contaminants and steroid hormones. Environmental Sciences, vol. 14, n° 6 (2007) 331-347
- Gutiérrez S, Fernández C, Escher BI, Tarazona JV (2008). A new hazard index of complex mixtures integrates bioconcentration and toxicity to refine the environmental risk assessment of effluents. Environ Int. 34(6), 773-81
- Haddad S, Charest-Tardif G, Tardif R, Krishnan K. (2000) Validation of a physiological modelling framework for simulating toxicokinetics of chemicals in mixtures. Toxicol Appl Pharmacol 2000; 167:199-209.
- Haigler BE, Pettigrew CA, Spain JC (1992). Biodegradation of Mixtures of Substituted Benzenes by Pseudomonasp. Strain JS150. Appl Environ Microbiol. 58(7), 2237-2244
- Hass U, Scholze M, Christiansen S, Dalgaard M, Vinggaard AM, Axelstad M, Metzdorff SB, Kortenkamp A (2007). Combined exposure to anti-androgens exacerbates disruption of sexual differentiation in the rat. Environment Health Perspect 115 (Suppl 1), 122-128
- Haws LC, Su SH, Harris M, Devito MJ, Walker NJ, Farland WH, Finley B, Birnbaum LS (2006). Development of a refined database of mammalian relative potency estimates for dioxin-like compounds. Toxicol Sci. 89, 4-30
- Heath E, Brown WA, Jensen SR, Bratty MP (2006). Biodegradation of chlorinated alkanes and their commercial mixtures by Pseudomonasp. strain 273. J Ind Microbiol Biotechnol 33(3),197-207
- Hermens J, Canton H, Janssen P, De Jong R (1984). Quantitative structure-activity relationships and toxicity studies of mixtures of chemicals with anaesthetic potency: acute lethal and sublethal toxicity to Daphnia magna. Aquatic Toxicology 5, 143-154
- Hermens J, Leeuwangh P, Musch A (1985). Joint toxicity of mixtures of groups of organic aquatic pollutants to the guppy (Poecilia reticulata). Ecotoxicol Environ Safety 9, 321-326
- Herrman JL and Younes M (1999). Background to the ADI/TDI/PTWI . Regul Toxicol Pharmacol 30, S109-S113
- Hulzebos EM, Posthumus R (2003). (Q)SARSS: gatekeepers against risk on chemicals? SAR QSAR Environ Res 14(4), 285–316
- IFRA (International Fragrance Association (2011) Considerations for Mixtures of Fragrance Ingredients - Update 2011 Final.doc September 2, 2011 1/9
- Jobling S, Burn RW, Thorpe K, Williams R, Tyler Ch (2009). Statistical Modeling Suggest that Antiandrogens in Effluents from Wastewater Treatment Works Contribute to Widespread Sexual Disruption in Fish Living in English Rivers. Environment Health Perspect 117(5), 797-802
- JRC (2010). SCIENTIFIC REPORT submitted to EFSA. Applicability of QSAR analysis to the evaluation of the toxicological relevance of metabolites and degradates of pesticide active substances for dietary risk assessment. Prepared by Computational Toxicology Group, Institute for Health & Consumer Protection, European Commission - Joint Research Centre, Ispra, Italy (Question No EFSA-Q-2009-01076. Accepted for Publication on 5 May 2010)

- Junghans M, Backhaus T, Faust M, Scholze M, Grimme LH (2006). Application and validation of approaches for the predictive hazard assessment of realistic pesticide mixtures. *Aquat. Toxicol.* 76, 93–110
- Kacham R, Karanth S, Baireddy P, Liu J, Pope C (2006). Interactive toxicity of chlorpyrifos and parathion in neonatal rats: role of esterases in exposure sequence-dependent toxicity. *Toxicol Appl Pharmacol.* 210(1-2), 142-149
- Karanth S, Liu J, Olivier K Jr, Pope C (2004). Interactive toxicity of the organophosphorus insecticides chlorpyrifos and methyl parathion in adult rats. *Toxicol Appl Pharmacol.* 196(2), 183-190
- Karanth S, Olivier K Jr, Liu J, Pope C (2001). In vivo interaction between chlorpyrifos and parathion in adult rats: Sequence of administration can markedly influence toxic outcome. *Toxicol Appl Pharmacol.* 177(3), 247-255
- Kavlock R., Dix D. (2010). Computational toxicology as implemented by the US EPA: providing high throughput decision support tools for screening and assessing chemical exposure, hazard and risk. *Journal of Toxicology and Environmental Health, Part B*, 13:197–217, 2010
- King DJ, Lyne RL, Girling A, Peterson DR, Stephenson R, Short D (1996). Environmental risk assessment of petroleum substances: The hydrocarbon block method. CONCAWE 96/52. Conservation of Clean Air and Water in Europe, Brussels, Belgium
- Knightes CD, Peters CA (2006). Multisubstrate biodegradation kinetics for binary and complex mixtures of polycyclic aromatic hydrocarbons. *Environ Toxicol Chem.* 25(7), 1746-56
- Korsak Z, Sokal J, Dedyk A, Tomas T, Jedrychowski R (1988). Toxic effects of combined exposure to toluene and xylene in animals. I. Acute inhalation study. *P J Occup Med* 1, 45-50
- Korsak Z, Sokal JA, Górny R. (1992) Toxic effects of combined exposure to toluene and m-xylene in animals. III. Subchronic inhalation study. *Pol J Occup Med Environ Health*; 5:27-33
- Kortenkamp A, Backhaus T, Faust M (2009). State of the Art on Mixture Toxicity. Report. Available at: http://ec.europa.eu/environment/chemicals/pdf/report_Mixture%20toxicity.pdf . Last accessed 15 April 2011
- Kortenkamp A., Faust M. (2010). Combined exposures to anti-androgenic chemicals: steps towards cumulative risk assessment. *International Journal of Andrology* 33 (2010), 463–474
- Kortenkamp A., Hass U. (2009). Expert workshop on combination effects of chemicals, 28-30. January 2009, Hornbæk, Denmark. Organized under the auspices of the Danish Ministry of the Environment and the Danish Environmental Protection Agency. Workshop Report. June 2009
- Laetz CA, Baldwin DH, Collier TK, Hebert V, Stark JD, et al. (2008) The Synergistic Toxicity of Pesticide Mixtures: Implications for Risk Assessment and the Conservation of Endangered Pacific Salmon. *Environ Health Perspect* 117(3): 117:348-53
- Lepper P (2005) Manual on the Methodological Framework to Derive Environmental Quality Standards for Priority Substances in accordance with Article 16 of the Water Framework Directive (2000/60/EC). Fraunhofer-Institute Molecular Biology and Ecology Schmallenberg, Germany. 15 September 2005
- Leslie HA., ter Laak TL., BUSSEER FM., Kraak MS., Hermens JLM., (2002) Bioconcentration of Organic Chemicals: Is a Solid-Phase Microextraction Fiber a Good Surrogate for Biota? *Environ. Sci. Technol.* 2002, 36, 5399-5404

- Levy JI (2008). Is epidemiology the key to cumulative Risk assessment? *Risk Analysis* 28, 1507-1513
- Loewe S and Muischnek H (1926). Über Kombinationswirkungen. *Naunyn-Schmiedebergs Arch Exp Pathol Pharmacol* 114, 313-326
- Mauderly JL, Samet JM (2009). Is there evidence for synergy among air pollutants in causing health effects? *Env Health Perspect* 117, 1-6
- Meek ME, Boobis AR, Crofton KM, Heinemeyer G, Raaij MV, Vickers C (2011), Risk assessment of combined exposure to multiple chemicals: A WHO/IPCS framework, *Regulatory Toxicology and Pharmacology* 60 (2011) S1-S14
- Meyer SA, Kim TW, Moser GJ, Monteiro-Riviere NA, Smart RC (1994). Synergistic interaction between the nonphorbol ester-type promoter mirex and 12-o-tetradecanoylphorbol-13-acetate in mouse skin tumor promotion. *Carcinogenesis* 15, 47-52
- Moser VC, Casey M, Hamm A, Cater WH Jr, Simmons JE, Gennings C (2005). Neurotoxicological and statistical analyses of a mixture of five organophosphorus pesticides using a ray design. *Toxicol Sci* 86, 101-115
- Moser VC, Simmons JE, Gennings C (2006). Neurotoxicological interactions of a five-pesticide mixture in preweanling rats. *Toxicol Sci* 92, 235-245
- Nesnow S, Mass MJ, Ross JA, Galati ANJ, Lambert GR, Gennings C, Carter Jr WH, Stoner GD (1998). Lung Tumorigenic Interactions in Strain A/J Mice of Five Environmental Polycyclic Aromatic Hydrocarbons. *Environ Health Perspect* 1998; 106 Suppl 6:1337-46.
- NTP-CERHR-BPA-07 (2007) NTP-CERHR Expert Panel Report on the Reproductive and Developmental Toxicity of Bisphenol A, Center for the Evaluation of Risks to Human Reproduction, November 26, 2007
- OECD (2009). Guidance document for using the OECD (Q)SAR Application Toolbox to develop chemical categories according to the OECD Guidance on grouping of chemicals. ENV/JM/MONO(2009)5, Series on Testing and Assessment No. 102
- OECD (2011) WHO OECD ILSI/HESI International Workshop on Risk Assessment of Combined Exposures to Multiple Chemicals. Paris, France, OECD Environment Directorate. OECD Environment, Health and Safety Publications. Series on Testing and Assessment.
- Parkerton TF., Stone MA., Letinski DJ. (2000) Assessing the aquatic toxicity of complex hydrocarbon mixtures using solid phase microextraction. *Toxicology Letters* 112-113 (2000) 273-282
- Payne J, Scholze M, Kortenkamp A (2001). Mixtures of four organochlorines enhance human breast cancer cell proliferation. *Environ Health Perspect.* 109, 391-197
- Plackett RL and Hewlett PS (1952). Quantal responses to mixtures of poisons. *J Royal Stat Soc Ser B.* 14, 141-154
- Plackett RL, Hewlett PS. (1948) Statistical aspects of the independent joint action of poisons, particularly insecticides; the toxicity of a mixture of poisons. *Ann Appl Biol* 35:347-58.
- Posthuma L., de Zwart D. (2006). Predicted effects of toxicant mixtures are confirmed by changes in fish species assemblages in Ohio, USA, rivers. *Environmental Toxicology and Chemistry*, Vol. 25, No. 4, pp. 1094-1105, 2006
- Price P and Han X (2011). Determining the need for cumulative risk assessments: When are chemical-by- chemical approaches insufficient and by how much? SETAC Special Symposium, Brussels, February 2011

- Price PS, Hollnagel HM, Zahik JM (2009). Characterizing the noncancer toxicity of mixtures using concepts from the TTC and quantitative models of uncertainty in mixture toxicity. *Risk Analysis* 29(11), 1534-1548
- Price PS., Chaisson CF. (2005). A conceptual framework for modeling aggregate and cumulative exposures to chemicals. *Journal of Exposure Analysis and Environmental Epidemiology* (2005) 15, 473-481
- Price PS., Han X. (2011). Maximum Cumulative Ratio (MCR) as a Tool for Assessing the Value of Performing a Cumulative Risk Assessment. *Int. J. Environ. Res. Public Health* 2011, 8, 2212-2225
- Price PS., Young JS., Chaisson CF. (2001). Assessing Aggregate and Cumulative Pesticide Risks Using a Probabilistic Model. *Ann. occup. Hyg.*, Vol. 45, No. 1001, pp. S131-S142, 2001
- Rajapakse N, Silva E, Scholze M, Kortenkamp A (2004). Deviation from additivity with estrogenic mixtures containing 4-nonylphenol and 4-tert-octylphenol detected in the E-SCREEN assay", *Environmental Science & Technology* 38(23), 6343-6352
- Reardon KF, Mosteller DC, Rogers JB, DuTeau NM, Kim KH (2002). Biodegradation Kinetics of Aromatic Hydrocarbon Mixtures by Pure and Mixed Bacterial Cultures. *Environ Health Perspect.* 110(suppl 6), 1005-1011
- Rider CV., LeBlanc GA. (2005) An Integrated Addition and Interaction Model for Assessing Toxicity of Chemical Mixtures. *Toxicological Sciences* 87(2), 520-528 (2005)
- Rusyn I and Daston GP (2010). Computational Toxicology: Realizing the promise of the toxicity testing in the 21st century. *Environ Health Perspect* 118, 1047-1050
- Schaafsma G., Kroese ER., Tielemans ELJP., van de Sandt JJM., van Leeuwen CJ. (2009) non-testing approaches and the urgent need for a change in mind set. *Regulatory Toxicology and Pharmacology* 53 (2009) 70-80
- SCHER (2007), Opinion on risk assessment on indoor air quality, 29 May 2007. http://ec.europa.eu/health/archive/ph_risk/committees/04_scher/docs/scher_o_055.pdf (Accessed on 14 Dec. 2011)
- Seed J., Carney EW., Corley RA., Crofton KM., DeSesso JM., Foster PMD., Kavlock R., Kimmel G., Klaunig J., Meek ME., Priston RJ., Slikker W Jr., Tabacova S., Williams GM., Wiltse J., Zoeller RT., Fenner-Crisp P., Patton DE. (2005). Overview: Using Mode of Action and Life Stage Information to Evaluate the Human Relevance of Animal Toxicity Data. *Critical Reviews in Toxicology*, 35:663-672, 2005
- Steinberg P., Prutner W., Nicken P., Hamscher G. (2011). H295R cells: an in vitro model for the risk assessment of single fungicides and mixtures of them modulating estrone biosynthesis. *Altex* 28, Special Issue, Montreal 2011; P 54
- Stewart AG and Carter J (2009). Towards the development of a multidisciplinary understanding of the effects of toxic chemical mixtures on health. *Environ Geochem Health* 31, 239-251
- Struijs J., van der Grinten E., Aldenberg T. (2009) Toxic pressure in the Dutch delta measured with bioassays. Trends over the years 2000 - 2009. RIVM Report 607013013/2010. RIVM, The Netherlands
- Sundberg H., Hanson M., Liewenborg B., Zebühr Y., Broman D., Balk L. (2007). Dredging Associated Effects: Maternally Transferred Pollutants and DNA Adducts in Feral Fish. *Environ. Sci. Technol.* 2007, 41, 2972-2977
- Sundberg H., Hanson M., Liewenborg B., Zebühr Y., Broman D., Balk L. (2007). Supporting information. Dredging Associated Effects: Maternally Transferred Pollutants and DNA Adducts in Feral Fish.

- Tichý M, Borek-Dohalský V, Rucki M, Reitmajer J, Feltl L (2002). Risk assessment of mixtures: possibility of prediction of interaction between chemicals. *Int Arch Occup Environ Health* 75 (Suppl), S133-S136
- Timchalk C and Poet TS (2008). Development of a physiologically based pharmacokinetic and pharmacodynamic model to determine dosimetry and cholinesterase inhibition for a binary mixture of chlorpyrifos and diazinon in the rat. *Neurotoxicology* 29(3), 428-443
- Tornero-Velez R., Egeghy PP, and Cohen Hubal EA. (2011) Biogeographical Analysis of Chemical Co-Occurrence Data to Identify Priorities for Mixtures Research. *Risk Analysis*, DOI: 10.1111/j.1539-6924.2011.01658.x
- Tornero-Velez R., Egeghy PP., Cohen Hubal EA. (2011). Biogeographical Analysis of Chemical Co-Occurrence Data to Identify Priorities for Mixtures Research. *Risk Analysis* DOI: 10.1111/j.1539-6924.2011.01658.
- Traas TP., van de Meent D., Posthuma L., Hamers T., Kater BJ., de Zwart D., Aldenberg T. (2002) The potentially affected fraction as a measure of ecological risk. *Book Species Sensitivity Distributions in ecotoxicology*. pp. 316-343
- USEPA (2002). Supplemental Guidance for Conducting Health Risk Assessment of Chemical Mixtures as a Supplement to the EPA's Guidelines for the Health Risk Assessment of Chemical Mixtures (USEPA, 1986)
- Vainio H and Boffetta P (1994). Mechanisms of the combined effect of asbestos and smoking in the etiology of lung cancer. *Scand J Work Environ health* 20, 235-242
- Van den Berg M, Birnbaum LS, Denison M, De Vito M, Farland W, Feeley M, Fiedler H, Hakansson H, Hanberg A, Haws L, Rose M, Safe S, Schrenk D, Tohyama C, Tritschler A, Tuomisto A, Tysklind M, Walker N, Peterson RE (2006). The 2005 World Health Organization reevaluation of human and mammalian toxic equivalency factors for dioxins and dioxin-like compounds. *Toxicol Sci.* 93, 223-241
- Van Leeuwen K, Schultz T.W., Henry T., Diderich B., Veith GD, 2009. Using chemical categories to fill data gaps in hazard assessment. *SAR QSAR Environ. Res.* 20, 207-220
- Van Wijngaarden RPA., Cuppen JGM., Arts GHP., Crum SJH., van den Hoorn MW, van den Brink PJ., Brock TCM. (2004). Aquatic risk assessment of a realistic exposure to pesticides used in bulb crops: a microcosm study. *Environmental Toxicology and Chemistry*, Vol. 23, No. 6, pp. 1479-1498, 2004
- Verbruggen EMJ, Beek M., Pijnenburg J., Traas TP. (2008) Ecotoxicological environmental risk limits for total petroleum hydrocarbons on the basis of internal lipid concentrations. *Environmental Toxicology and Chemistry*, Vol. 27, No. 12, pp. 2436-2448, 2008
- Verbruggen EMJ., Vaes WHJ., Parkerton TF., Hermens JPL. (2000) Polyacrylate-Coated SPME Fibers as a Tool To Simulate Body Residues and Target Concentrations of Complex Organic Mixtures for Estimation of Baseline Toxicity. *Environ. Sci. Technol.* 2000, 34, 324-331
- Verbruggen EMJ., van den Brink PJ. (2010). Review of recent literature concerning mixture toxicity of pesticides to aquatic organisms. Report 601400001/2010. RIVM, the Netherlands.
- Verbruggen EMJ., van Loon WMGM., Tonkes M., van Duijn P., Seinen W., Hermens JLM. (1999) Biomimetic Extraction as a Tool To Identify Chemicals with High Bioconcentration Potential: An Illustration by Two Fragrances in Sewage Treatment Plant Effluents and Surface Waters. *Environ. Sci. Technol.* 1999, 33, 801-806
- Verro R, Finizio A, Otto S, Vighi M (2009). Predicting pesticide environmental risk in intensive agricultural areas. II: screening level risk assessment of complex mixtures in surface waters. *Environ Sci Technol.* 43, 530-537

- Viaene M., Vermeir G., van Larebeke N. (2007). Invloed van industriële chemicaliën op de neurologische en neuropsychologische ontwikkeling van jonge kinderen.
- Vighi M (2006) Unpublished data from a report of the BEAM project. Contract EVK1-CT999-00012
- VKM (2008). Norwegian Scientific Committee for Food Safety. Combined toxic effects of multiple chemical exposures. Report
- Walker NJ, Crockett PW, Nyska A, Brix AE, Jokinen MP, Sells DM, Hailey JR, Easterling M, Haseman JK, Yin M et al. (2005). Dose-additive carcinogenicity of a defined mixture of "dioxin-like" compounds. *Environ Health Perspect* 113, 43-48
- Walter H, Consolaro F, Gramatica P, Scholze M, Altenburger R (2002). Mixture toxicity of priority pollutants at no observed effect concentrations (NOECs). *Ecotoxicol.* 11, 299-310
- WHO (2009a). World Health Organization. Assessment of Combined Exposures to Multiple Chemicals: Report of a WHO/IPCS International Workshop
- WHO (2009b). World Health Organization. Harmonization Project. DRAFT Document for Public and Peer Review. Risk Assessment of Combined Exposures to Multiple Chemicals: A WHO/IPCS Framework
- Wolansky MJ, Gennings C, DeVito MJ, Crofton KM. (2009). Evidence for dose-additive effects of pyrethroids on motor activity in rats. *Environ Health Perspect* 117:1563-1570.
- Wraith D and Mengersen K (2007). Assessing the combined effect of asbestos exposure and smoking on lung cancer: a Bayesian approach. *Stat Med* 26, 1150-69.
- Zeliger HI. (2003) Toxic effects of chemical mixtures. *Arch Environ Health* 2003; 58(1):23-29
- Zeliger HI. (2008) Human toxicology of chemical mixtures. William Andrew, Norwich, New York. ISBN: 978-0-8155-1589-0
- Zeliger HI. Respiratory toxicity of chemical mixtures: predicting low level responses. Zeliger Chemical, Environmental & Toxicological Research & Services. West Charlton, New York. hiz@zeliger.com