

## **Bibliography of the Sources Examined for the Opinion on Nitrosamines and Secondary amines in Cosmetic Products**

- Arai M, Aoki Y, Nakanishi K, Miyata Y, Mori T, Iyo N (1979). Long-term experiment of maximal non-carcinogenic dose of dimethylnitrosamine for carcinogenesis in rats. *Gann* 70: 549-558.
- Bachrach, U. The early history of polyamine research. *Plant Physiology and Biochemistry* 48, (2010) 490-495
- Bae, J.-Y., et al., The Nitrosation of Hexetidine and Hexidine: Characterization of the Major Nitrosamine from Common Antimicrobial Agents. *Chem. Res. Toxicol.*, 1994. 7(6): p. 868-876.
- Bandeira, C.M. et al. Bioactive amines in fresh, canned and dried sweet corn, embryo and endosperm and germinated corn. *Food Chemistry* 131 (2012), 1355-1359
- Bayerisches Landesamt für Gesundheit und Lebensmittelsicherheit, Jahresberichte 2007-2010, <http://www.lgl.bayern.de/publikationen/jahresberichte.htm>
- Berger MR, Schmähl D, Edler L (1990) Implications of the carcinogenic hazard of low doses of three hepatocarcinogenic N-nitrosamines. *Jpn J Cancer Res* 81: 598-606.
- Berger MR, Schmähl D, Edler L (1990) Implications of the carcinogenic hazard of low doses of three hepatocarcinogenic N-nitrosamines. *Jpn J Cancer Res* 81: 598-606.
- Berger MR, Schmähl D, Zerban H (1987). Combination experiments with very low doses of three genotoxic N-nitrosamines with similar organotropic carcinogenicity in rats. *Carcinogenesis* 8: 1635-1643.
- Berger MR, Schmähl D, Zerban H (1987). Combination experiments with very low doses of three genotoxic N-nitrosamines with similar organotropic carcinogenicity in rats. *Carcinogenesis* 8: 1635-1643.
- Berwanger, A. et al. Polyamines: Naturally occurring small molecule modulators of electrostatic protein-protein interactions. *J. Inorganic Chemistry* 104 (2010) 118-125
- Bills, D.D., et al., Potential precursors of N-nitrosopyrrolidine in bacon and other fried foods. *Journal of Agricultural and Food Chemistry*, 1973. 21(5): p. 876-7.
- Bills, D.D., et al., Potential precursors of N-nitrosopyrrolidine in bacon and other fried foods. *J Agric Food Chem*, 1973. 21(5): p. 876-7.
- Bouffard Fita F. Statement from Sinnoqa regarding opinion on nitrosamines and secondary amines in cosmetic products SCCS/1458/11 (Scientific Committee on Consumer Safety). 17 January 2012
- Bull RJ, Reckhow DA, Li X, Humpage AR, Joll C, Hruedy SE. Potential carcinogenic hazards of non-regulated disinfection by-products: Haloquinones, halo-cyclopentene and cyclohexene derivatives, N-halamines, halonitriles, and heterocyclic amines. *Toxicology*. 2011 Aug 15;286(1-3):1-19. Epub 2011 May 14.
- Challis, B.C. and B.F.L. Li, Formation of N-nitrosamines and N-nitramines by photolysis. *IARC Sci. Publ.*, 1982. 41(N-Nitroso Compd: Occurrence Biol. Eff.): p. 31-40.
- Challis, B.C. and D.E.G. Shuker, Rapid nitrosation of amines in aqueous alkaline solutions by beta -substituted alkyl nitrites. *J. Chem. Soc., Chem. Commun.*, 1979(7): p. 315-16.
- Challis, B.C. and J.R. Outram, The chemistry of nitroso compounds. Part 15. Formation of N-nitrosamines in solution from gaseous nitric oxide in the presence of iodine. *J. Chem. Soc., Perkin Trans. 1*, 1979(11): p. 2768-75.

- Challis, B.C. and S.A. Kyrtopoulos, Rapid formation of carcinogenic N-nitrosamines in aqueous alkaline solutions. Br. J. Cancer, 1977. 35(5): p. 693-6.
- Challis, B.C. and S.A. Kyrtopoulos, The chemistry of nitroso compounds. Part 12. The mechanism of nitrosation and nitration of aqueous piperidine by gaseous dinitrogen tetraoxide and dinitrogen trioxide in aqueous alkaline solutions. Evidence for the existence of molecular isomers of dinitrogen tetraoxide and dinitrogen trioxide. J. Chem. Soc., Perkin Trans. 2, 1978(12): p. 1296-302.
- Challis, B.C. and S.A. Kyrtopoulos, The chemistry of nitroso-compounds. Part 11. Nitrosation of amines by the two-phase interaction of amines in solution with gaseous oxides of nitrogen. J. Chem. Soc., Perkin Trans. 1, 1979(2): p. 299-304.
- Challis, B.C., et al., Formation of N-nitrosamines and N-nitramines by gaseous nitrogen dioxide. Acta Cient. Compostelana, 1982. 19(1-2): p. 153-66.
- Challis, B.C., et al., Rapid formation of N-nitrosamines from nitrogen oxides under neutral and alkaline conditions. IARC Sci. Publ., 1978. 19(Environ. Aspects N-Nitroso Compd.): p. 127-42.
- Challis, B.C., J.R. Outram, and D.E.G. Shuker, New pathways for the rapid formation of N-nitrosamines under neutral and alkaline conditions. IARC Sci. Publ., 1980. 31(N-Nitroso Compd.: Anal., Form. Occurrence): p. 43-58.
- Chemisches und Veterinäruntersuchungsamt (CVUA) Karlsruhe, Jahresbericht 2006, [http://www.ua-bw.de/pub/beitrag.asp?subid=2&ID=991&Thema\\_ID=1&lang=DE](http://www.ua-bw.de/pub/beitrag.asp?subid=2&ID=991&Thema_ID=1&lang=DE)
- Chemisches und Veterinäruntersuchungsamt (CVUA) Karlsruhe, Jahresbericht 2005, [http://www.ua-bw.de/pub/beitrag.asp?subid=2&ID=992&Thema\\_ID=1&lang=DE](http://www.ua-bw.de/pub/beitrag.asp?subid=2&ID=992&Thema_ID=1&lang=DE)
- Chemisches und Veterinäruntersuchungsamt (CVUA) Stuttgart, Jahresbericht 2008, <http://jahresbericht.cvuas.de/docs/jb2008.pdf>
- Chemisches und Veterinäruntersuchungsamt (CVUA) Stuttgart, Jahresbericht 2007, <http://jahresbericht.cvuas.de/docs/jb2007.pdf>
- Chemisches und Veterinäruntersuchungsamt (CVUA) Stuttgart, Jahresbericht 2006, [http://cvuas.untersuchungsaemter-bw.de/pdf/cvuas\\_jahresbericht2006.pdf](http://cvuas.untersuchungsaemter-bw.de/pdf/cvuas_jahresbericht2006.pdf)
- Chemisches und Veterinäruntersuchungsamt (CVUA) Stuttgart, Jahresbericht 2005, [http://cvuas.untersuchungsaemter-bw.de/pdf/cvuas\\_jahresbericht2005.pdf](http://cvuas.untersuchungsaemter-bw.de/pdf/cvuas_jahresbericht2005.pdf)
- Chung FL, Tanaka T, Hecht SS (1986). Induction of liver tumors in F344 rats by crotonaldehyde. Cancer Research 46: 1285-1289.
- Colipa, Technical Guidance Document on Minimising and Determining Nitrosamines in Cosmetics. 2009, Colipa, The European Cosmetics Association. p. 43.
- Conrad, J., K.H. Schlemmer, and G. Eisenbrand, Nitrosation of Bromhexine, in The Significance of N-Nitrosation of Drugs, G. Eisenbrand, G. Bozler, and H.v. Nicolai, Editors. 1990, Gustav Fischer Verlag: Stuttgart. p. 181-196.
- Dadakova, E. et al. Concentration of biologically active polyamines in rabbit meat, liver and kidney after slaughter and their changes during meat storage and cooking. Meat Science 90 (2012) 796-800
- Dadakova, E. et al. Concentration of biologically active polyamines in meat and liver of sheep and lambs after slaughter and their changes in mutton during storage and cooking. Meat Science 87 (2011) 119-124
- De Mey, E. et al. Dabsyl derivatisation as an alternative for dansylation in the detection of biogenic amines in fermented meat products by reversed phase high performance liquid chromatography. Food Chemistry (2012), 130 (4), 1017-1023

Drabik-Markiewicz, G. et al. Influence of putrescine, cadaverine, spermidine or spermine on the formation of N-nitrosamine in heated cured pork meat. Food Chemistry 126 (2011) 1530-1545

Dybing E., Sanner T., Roelfzema H., Kroese D. and Tennant R.W. T25: A simplified carcinogenic potency index: Description of the system and study of correlations between carcinogenic potency and species/site specificity and mutagenicity. Pharmacology and Toxicology 80, 272-279 (1997).

ECHA (2008). Guidance on information requirements and chemical safety assessment; Chapter R.7c: Endpoint specific guidance, European Chemicals Agency, 2008

EFSA (2011) Use of BMDS and PROAST software packages by EFSA Scientific Panels and Units for applying the Benchmark Dose (BMD) approach in risk assessment. <http://www.efsa.europa.eu/en/supporting/doc/s113e.pdf>

EFSA Guidance of the Scientific Committee on Use of the benchmark dose approach in risk assessment. <http://www.efsa.europa.eu/en/efsjournal/pub/1150.htm>

Eisenbrand, G., Blankart M., Sommer H., Weber B., N-Nitrosoalkanolamines in cosmetics. In: Relevance to human cancer of N-nitroso compounds, tobacco smoke and mycotoxins, I.K. O'Neill, J. Chen and H. Bartsch, (eds.), 1991, IARC Sci. Publ. 105, pp. 238–241.

Eisenbrand, G., Endogenous nitrosation - findings and problems, in The Significance of N-Nitrosation of Drugs, G. Eisenbrand, G. Bozler, and H.v. Nicolai, Editors. 1990, Gustav Fischer Verlag: Stuttgart. p. 47-68.

Ellen, G., J. Freudenthal, and P.L. Schuller, N-Nitroso-1,3-oxazolidine, a carcinogenic compound detected as an impurity in a commercial cutting fluid. Recl.: J. R. Neth. Chem. Soc., 1982. 101(10): p. 357-9.

Fitzpatrick RB CPDB: Carcinogenic Potency Database. Med Ref Serv Q. 2008 Fall; 27(3):303-11.

Fuchs, T. et al. Content of polyamines in by-products of slaughter pigs. Meat Science 83 (2009) 161-164

Gewehr, C et al. Contribution of peripheral vanilloid receptor to the nociception induced by injection of spermine in mice. Pharmacology, Biochemistry and Behavior 99 (2011) 775-781

Gowenlock, B., et al., Nitrosative dealkylation of some symmetrical tertiary amines. J. Chem. Soc., Perkin Trans. 2, 1979: p. 1110-1114.

Gray R, Peto R, Brantom P, Grasso P (1991). Chronic nitrosamine ingestion in 1040 rodents: The effect of the choice of nitrosamine, the species studied, and the age of starting exposure. Cancer Research 51: 6470-6491.

Gupta, N. et al. Chronic difluoromethylornithine treatment impairs spatial learning and memory in rats. Pharmacology, Biochemistry and Behavior 100 (2012) 464-473

Habs M, Schmähl (1980). Synergistic effect of N-nitroso compounds in experimental long-term carcinogenesis studies. Oncology 37: 259-265.

Hecht SS, Lijinsky W, Kovatch RM, Chung FL, Saavedra JE (1989). Comparative tumorigenicity of N-nitroso-2-hydroxymorpholine, N-nitrosodiethanolamine and N-nitrosomorpholine in A/J mice and F344 rats. Carcinogenesis 10: 1475 – 1477.

Helguera AM, Pérez-Machado G, Cordeiro MN, Combes RD Quantitative structure-activity relationship modelling of the carcinogenic risk of nitroso compounds using regression analysis and the TOPS-MODE approach. SAR QSAR Environ Res. 2010 Apr; 21(3-4):277-304.

- Hernandez LG, Slob W, van Steeg H, van Bentham J. Can carcinogenic potency be predicted from in vivo genotoxicity data? A meta-analysis of historical data. Environ Mol Mutagen 52: 518-528, 2011.
- Hildrum, K.I. and R.A. Scanlan, Factors influencing the rate of formation of volatile N-nitrosamines during the nitrosation of spermidine. J. Agric. Food Chem., 1977. 25(2): p. 255-7.
- Hildrum, K.I., R.A. Scanlan, and L.M. Libbey, Formation of volatile, hydroxylated, and chlorinated N-nitrosamines during the nitrosation of spermidine 3-hydrochloride. J Agric Food Chem, 1977. 25(2): p. 252-5.
- Hildrum, K.I., R.A. Scanlan, and L.M. Libbey, Identification of gamma -butenyl-(beta -propenyl)nitrosamine, the principal volatile nitrosamine formed in the nitrosation of spermidine or spermine. J. Agric. Food Chem., 1975. 23(1): p. 34-7.
- Hildrum, K.I., R.A. Scanlan, and L.M. Libbey, Nitrosamines from the nitrosation of spermidine and spermine. IARC Sci Publ, 1976(14): p. 205-14.
- Hoos A, Habst M, Schmähle D (!985). Comparison of liver tumor frequencies after intermittent oral administration of different doses of N-nitrosopyrrolidine in Sprague-Dawley rats. Cancer Letters 26: 77-82.
- Hotchkiss, J.H., et al., Mutagenicity of nitrosamines formed from nitrosation of spermidine. Mutat Res, 1979. 68(3): p. 195-9.
- Hotchkiss, J.H., R.A. Scanlan, and L.M. Libbey, Formation of bis(hydroxyalkyl)-N-nitrosamines as products of the nitrosation of spermidine. J. Agric. Food Chem., 1977. 25(5): p. 1183-9.
- Huxel, E.T., R.A. Scanlan, and L.M. Libbey, Formation of N-nitrosopyrrolidine from pyrrolidine ring containing compounds at elevated temperatures. Journal of Agricultural and Food Chemistry, 1974. 22(4): p. 698-700.
- Ikeda, K et al. Analysis of nitrosamines in cosmetics. J. Soc. Cosmet. Chem. 41, 283-333 (1990)
- Janne, J. et al. Polyamines and polyamine-metabolizing enzyme activities in human semen. Clinica Chimica Acta, 48 (1973) 393-401
- Janzowski, C., Hemm, I., Eisenbrand, G., in Wichmann, Schlipkötter, Füllgraf, eds, Handbuch Umweltmed.11/00,Org.Verb./N-Nitrosamine IV-4.1-30, 2000°
- Jones, A.R., W. Lijinsky, and G.M. Singer, Steric effects in the nitrosation of piperidines. Cancer Res., 1974. 34(5): p. 1079-81.
- Keefer, L.K. and P.P. Roller, N-nitrosation by nitrite ion in neutral and basic medium. Science, 1973. 181(4106): p. 1245-1247.
- Klaveness, J. 2012. Spermine and other polyamines in food and food products
- Klaveness, J. 2012. Spermine and other polyamines. Function and presence in humans
- Klaveness, J. 2012. Spermine comprising cosmetics. Potential for formation of nitrosamines
- Kofink, M. et al. determination of nitrosamines. Certificate n° 1266063-01 to 08. SGS. 2012
- Konishi Y, Kondo H, Ikeda T, Kawabata A, Shoji Y, Deda A (1978). Effect of dose on the carcinogenic activity of orally administrated N-bis(2-hydroxypropyl)nitrosamine in rats. GANN 69: 573-577.
- Kozova, M. et al. Changes in content of biologically active polyamines during beef loin storage and cooking. Meat Science 81 (2009) 607-611

- Kozova, M. et al. Contents of biologically active polyamines in chicken meat, liver, heart and skin after slaughter and their changes during meat storage and cooking. *Food Chemistry* 116 (2009) 418-425
- Krizek, M et al. Effect on low-dose irradiation on biogenic amines formation in vacuum-packed trout flesch. *Food Chemistry* 132 (2012) 367-372
- Larque, E. et al. Biological significance of dietary polyamines. *Nutrition* 23 (2007) 87-95
- Lijinsky W, Kovatch M, Riggs CV (1987) Carcinogenesis by nitrosodialkylamines and azoxyalkanes given by gavage to rats and hamsters. *Cancer Res.* 47: 3968-3972.
- Lijinsky W, Kovatch RM (1985). Induction of liver tumors in rats by nitrosodiethanolamine at low doses. *Carcinogenesis* 6: 1679 – 1681.
- Lijinsky W, Kovatch RM, Riggs CW, Walters PT (1988). Dose response study with N-nitrosomorpholine in drinking water of F-344 rats. *Cancer Research* 48: 2089-2095.
- Lijinsky W, Reuber MD (1981). Carcinogenic effect of nitrosopyrrolidine, nitrosopiperidine and nitrosohexamethyleneimine in Fischer rats. *Cancer Letters* 12: 99-103.
- Lijinsky W, Reuber MD (1982). Comparative carcinogenesis by nitrosomorpholines, nitrosooxazolidines and nitrosotetrahydrooxazine in rats. *Carcinogenesis* 3: 911-915.
- Lijinsky W, Reuber MD (1984). Carcinogenesis in rats by nitrosodimethylamine and other nitrosomethylalkylamines at low doses. *Cancer Letters* 22: 83-88
- Lijinsky W, Reuber MD, Riggs CW (1981). Dose response studies of carcinogenesis in rats by nitrosodiethylamine. *Cancer Research* 41: 4997-5003 (1981).
- Lijinsky W, Reuber MD, Riggs CW (1983). Carcinogenesis by combinations of nitroso compounds in rats. *Food Chem. Toxicol.* 21: 601 – 605.
- Lijinsky W, Saavedra JE, Reuber MD (1984). Carcinogenesis in rats by some hydroxylated acyclic nitrosamines. *Carcinogenesis* 5: 167-170.
- Lijinsky W, Taylor HW (1976). The effect of substituents on the carcinogenicity of N-nitrosopyrrolidine in Sprague-Dawley Rats. *Cancer Research* 36: 1988-1990.
- Loeppky, R.N. and J.Y. Bae, An aziridinium ion intermediate in the nitrosation of a hexetidine model. *Chem. Res. Toxicol.*, 1994. 7: p. 661-667.
- Loeppky, R.N. and T. Tomasik, Stereoelectronic Effects in Tertiary Amine Nitrosation: Nitrosative Cleavage vs. Aryl Ring Nitrosation. *J. Org. Chem.*, 1983. 48: p. 2751-2756.
- Loeppky, R.N., et al., Blocking nitrosamine formation: Understanding the chemistry of nitrosamine formation, in *Nitrosamines and Related N-Nitroso Compounds: Chemistry and Biochemistry*, R.N. Loeppky and C.J. Michejda, Editors. 1994, American Chemical Society: Washington, DC. p. 52-65.
- Loeppky, R.N., et al., Evidence for radical cations in linked mechanisms of N,N-dialkyl aromatic amine nitration and nitrosative dealkylation. *J. Am. Chem. Soc.*, 1998. 120(21): p. 5193-5202.
- Loeppky, R.N., et al., Nitrosamine formation from ternary nitrogen compounds. International Agency for Research in Cancer Scientific Publication, 1982. 41: p. 41-56.
- Loeppky, R.N., et al., Nitrosation of tertiary aromatic amines related to sunscreen ingredients, in *Relevance of Human Cancer of N-Nitroso Compounds, Tobacco Smoke, and Mycotoxins*, I.K. O'Neill, J. Chen, and H. Bartsch, Editors. 1991, International Agency for Research on Cancer: Lyon, Fr. p. 244-252.
- Loeppky, R.N., G. Shevlin, and L. Yu, Rapid nitrosamine formation from tertiary nitrogen compounds: An overview, in *Significance of N-Nitrosation of Drugs (Drug Dev.*

- Eval), G. Eisenbrand, G. Bolzer, and H.v. Nicolai, Editors. 1990, Gustav Fischer Verlag: New York.
- Loeppky, R.N., W. Tomaszik, and T.G. Millard, Ester-mediated nitrosamine formation from nitrite and secondary or tertiary amines. International Agency for Research in Cancer Scientific Publication, 1984. 57: p. 353-363.
- Maduagwu E.N. Nitrosation of quaternary ammonium compounds in vivo in the Wistar rat. *Xenobiotica* 15(12):1061-1068.
- Mah, J.-H., et al., Influence of curing and heating on formation of N-nitrosamines from biogenic amines in food model system using Korean traditional fermented fish product. *Food Sci. Biotechnol.*, 2005. 14(1): p. 168-170.
- Mayer, H.K. et al. A new ultra-pressure liquid chromatography method for the determination of biogenic amines in cheese. *Journal of Chromatography A*, 1217 (2010) 3251-3257
- McGregor D, Boobis A, Binaglia M, Botham P, Hoffstadt L, Hubbard S, Petry T, Riley A, Schwartz D, Hennes C. Guidance for the classification of carcinogens under the Globally Harmonised System of Classification and Labelling of Chemicals (GHS). *Crit Rev Toxicol*. 2010 Mar; 40(3):245-85.
- Meier KL, Bailer AJ, Portier CJ (1993). A measure of tumorigenic potency incorporating dose-response shape. *Biometrics*. 49: 917-926
- Michejda CJ, Kroeger-Koepke MB, Kovatch RM (1986). Carcinogenic effects of sequential administration of two nitrosamines in Fischer 344 rats. *Cancer Research* 46: 2252-2256.
- Mirvish, S.S., Blocking the formation of N-nitroso compounds with ascorbic acid in vitro and in vivo. *Ann. N. Y. Acad. Sci.*, 1975: p. 258.
- Mirvish, S.S., Formation of N-Nitroso compounds: Chemistry, kinetics and in vivo occurrence. *Toxicology and Applied Pharmacology*, 1975. 31: p. 325-351.
- Mirvish, S.S., Kinetics of dimethylamine nitrosation in relation to nitrosamine carcinogenesis. *J. Nat. Cancer Inst.*, 1970. 44(3): p. 633-639.
- Naila, A. et al. Biogenic amines and potential histamine-Forming bacteria in Rihaakuru (a cooked fish paste). *Food Chemistry* 128 (2011) 479-484
- Oefner, P.J. et al. High-performance liquid chromatographic determination of free polyamines in human seminal plasma. *Clinica Chimica Acta*, 205 (1992) 11-18
- Okun,J.D, Archer,M.C. Kinetics of nitrosamine formation in the presence of micelle-forming surfactants. *J.Natl.Cancer Inst.* 1977,58(2): 409-11)
- Papavergou, E.J. Biogenic amine levels in dry fermented sausages produced and sold in Greece. *Procedia Food Science* 1 (2011) 1126-1131
- Park, J.S. et al. Monitoring the contents of biogenic amines in fish and fish products consumed in Korea. *Food Control* 21 (2010) 1219-1226
- Park, M., F. Gu, and R.N. Loeppky, The synthesis of highly reactive multifunctional -epoxy- and -acetoxy-nitrosamines. *Tetrahedron Lett.*, 1998. 39: p. 1287-1290.
- Paz, E.A. et al. Polyamines in cancer. *Advances in Clinical Chemistry* 54 (2011) 45-70
- Perera, A.M.A., N-nitrosoamines. *Chromatogr. Sci. Ser.*, 2006. 93(Chromatographic Analysis of the Environment (Third Edition)): p. 419-452.
- Peto R, Gray R, Brantom P, Grasso P (1991). Effects on 4080 rats of chronic ingestion of N-Nitrosodiethylamine or N-Nitrosodimethylamine: A Detailed Dose-Response Study. *Cancer Res* 51: 6415-6451.

- Pineda, A et al. Preliminary evaluation of biogenic amines content in Chilean young varietal wines by HPLC. *Food Control* 23 (2012) 251-257
- Portier CJ, Hoel DG (1987). Issues concerning the estimation of the TD50. *Risk Anal.* 7: 437-447.
- Preussmann R, Habs M, Habs H, Schmähl D (1982). Carcinogenicity of N-nitrosodiethanolamine in rats at five different dose levels. *Cancer Res* 42: 5167 – 5171.
- Preussmann R, Schmähl D, Eisenbrand G (1977). Carcinogenicity of N-Nitrosopyrrolidine: Dose-Response Study in Rats. *Z. Krebsforsch.* 90: 161-166.
- Preussmann,R in The Significance of n-Nitrosation of drugs,1990,G.Fischer Verl.Stuttg.New York, Carcinogenicity and Structure-Activity relationships of N-Nitroso compounds,p3-17
- Ray, R. M. et al. Spermine, a molecular switch regulating EGFR integrin beta-3, Src, and FAK Scaffolding. *Cellular Signaling* 24 (2012) 933-42
- RIVM Rijksinstituut voor Volksgezondheid en Milieu, Markennis Cosmetica, Nitrosaminen (NDELA) en Zware Metalen in Decoratieve Cosmetica, Fact Sheet, September 2009
- RIVM Rijksinstituut voor Volksgezondheit en Millieu, Nitrosaminen (NDELA) in cosmetica, Fact sheet, März 2008
- Rostkowska, K. et al. Formation and Metabolism of N-Nitrosamines. *Polish Journal of Environmental Studies* Vol. 7, n° 6 (1998), 321-325
- Sagratini, G et al. Simultaneous determination of eight underivatised biogenic amines in fish by solid phase extraction and liquid chromatography-tandem mass spectrometry. *Food Chemistry* 132 (2012) 537-543
- Sanner T, Dybing E. Comparison of carcinogenic and in vivo genotoxic potency estimates. *Basic Clinical Pharm Toxicol*, 96: 131–139, 2005.
- Sanner T., Dybing E., Willems M.I. and Kroese E.D. A simple method for quantitative risk assessment of non-threshold carcinogens based on the dose descriptor T25. *Pharmacol Toxicol* 88: 331-341 (2001).
- Santiago-Silva, P. et al. Functional potential of tropical fruits with respect to free bioactive amines. *Food Research International* 44 (2011) 1264-1268
- SCCS's Notes of Guidance for the testing of Cosmetic Ingredients and their Safety evaluation, of 14 December 2010
- SCHER/SCCP/SCENIHR, 2009 - Scientific opinion on risk assessment methodologies and approaches for genotoxic and carcinogenic substances, adopted on the 19th plenary meeting of the SCCP of 21 January 2009.
- Schothorst, R.C., and Somers, H.H.J., Determination of N-nitrosodiethanolamine in cosmetic products by LC-MS-MS. *Anal. Bioanal. Chem.* 2005, 381, 681-685.
- Schothorst, R.C., and Stephany, R.W., Occurrence of N-nitrosodiethanolamine (NDELA) in cosmetics from the Dutch market. *Int. J. Cosmet. Sci.* 2001, 23, 109-114.
- Sedgwick, B., Nitrosated peptides and polyamines as endogenous mutagens in O6-alkylguanine-DNA alkyltransferase deficient cells. *Carcinogenesis*, 1997. 18(8): p. 1561-1567.
- Shukia, S. et al. Determination of biogenic amines in Korean traditional fermented soybean paste (Doenjang). *Food and Chemical Toxicology* 48 (2010) 1191-1195
- Skotvold T. 2012. Background, summary of reply and comments to the SCCS preliminary opinion on "Nitrosamines and Secondary Amines in Cosmetic Products". Scandinavian Dermal Innovations

- Smith, P.A.S. and R.N. Loeppky, Nitrosative cleavage of tertiary amines. *J. Am. Chem. Soc.*, 1967. 89(5): p. 1147-1157.
- Stephany, R.W., J. Freudenthal, and P.L. Schuller, N-Nitroso-5-methyl-1,3-oxazolidine identified as an impurity in a commercial cutting fluid. *Recueil des Travaux Chimiques des Pays-Ba*, 1978. 97(6): p. 177-8.
- Tang, T. et al. Monitoring the contents of biogenic amines in sufu by HPLC with SPE and pre-column derivatization. *Food Control* 22 (2011) 1203-1208
- Tasic, T. et al. Biogenic amines content in traditional dry fermented sausage Petrovák klobása as possible indicator of good manufacturing practice. *Food Control* 23 (2012) 107-112
- Terao K, Alkawa T, Kera K (1978). A synergistic effect of nitrosodimethylamine on sterigmatocystin carcinogenesis in rats. *Fd Cosmetic Toxicol* 16: 591-596.
- Teuten, E.L. and R.N. Loeppky, The mechanistic origin of regiochemical changes in the nitrosative N-dealkylation of N,N-dialkyl aromatic amines. *Organic & Biomolecular Chemistry*, 2005. 3(6): p. 1097-1108.
- Triki, M. et al. Optimisation of chromatographic procedure for determining biogenic amine concentrations in meat and meat products employing a cation-exchange column with a post-column system. *Food Chemistry* 130 (2012) 1066-1073
- Wei, F et al. Irradiated Chinese Rugao ham: Change in volatile N-nitrosamine, biogenic amine and residual nitrite during ripening and post-ripening. *Meat Science* 81 (2009) 451-455
- Williams, D.L.H., Nitrosation reactions in the chemistry of nitric oxide. 2004, Amsterdam: Elsevier. 268.
- Yongmei, L. et al. Biogenic amines in Chinese soy sauce. *Food Control* 20 (2009) 593-597
- Zarei, M. et al. Chemical and microbial properties of mahyaveh, a traditional Iranian fish sauce. *Food Control* 23 (2012) 511-514
- Zerban H, Preussmann R, Bannasch P (1988). Dose-time relationship of the development of preneoplastic liver lesions induced in rats with low doses of N-nitrosodiethanolamine. *Carcinogenesis* 9: 607 – 610.
- Zhai, H. et al. Biogenic amines in commercial fish and fish products sold in southern China. *Food Control* 25 (2012) 303-308