Addendum to the scientific opinion on o-Phenylphenol, Sodium o-phenylphenate and Potassium o-phenylphenate SCCS/1555/15



# Scientific Committee on Consumer Safety SCCS

# **ADDENDUM**

to the scientific opinion on o-Phenylphenol, Sodium ophenylphenate and Potassium o-phenylphenate (SCCS/1555/15)

**Here:** the use as preservative of

Sodium o-phenylphenate, Potassium o-phenylphenate, MEA o-Phenylphenate

(CAS n. 132-27-4, 13707-65-8, 84145-04-0)



The SCCS adopted this document at its plenary meeting on 21-22 February 2018

\_\_\_\_\_\_

#### **ACKNOWLEDGMENTS**

#### SCCS members

Dr U. Bernauer (

(Rapporteur)

Dr L. Bodin

Prof. Q. Chaudhry

Prof. P.J. Coenraads

Prof. M. Dusinska

Dr J. Ezendam

Dr E. Gaffet

Prof. C. L. Galli

Dr B. Granum

Prof. E. Panteri

Prof. V. Rogiers

Dr Ch. Rousselle

Dr M. Stepnik

Prof. T. Vanhaecke

Dr S. Wijnhoven

All Declarations of Working Group members are available on the following webpage: <a href="http://ec.europa.eu/health/scientific committees/experts/declarations/sccs">http://ec.europa.eu/health/scientific committees/experts/declarations/sccs</a> en.htm

In agreement with the mandating DG, there is no commenting period for this Addendum.

Keywords: SCCS, scientific opinion, preservative, Sodium o-phenylphenate, Potassium o-phenylphenate, MEA o-Phenylphenate, Regulation 1223/2009, CAS n. 132-27-4, 13707-65-8, 84145-04-0

Opinion to be cited as: SCCS (Scientific Committee on Consumer Safety), Addendum to the scientific opinion on the use as preservative of o-Phenylphenol, Sodium o-phenylphenate and Potassium o-phenylphenate- Here: the use as preservative of Sodium o-phenylphenate, Potassium o-phenylphenate, MEA o-Phenylphenate 21-22/02/2018, SCCS/1597/18

#### About the Scientific Committees

Two independent non-food Scientific Committees provide the Commission with the scientific advice it needs when preparing policy and proposals relating to consumer safety, public health and the environment. The Committees also draw the Commission's attention to the new or emerging problems which may pose an actual or potential threat.

They are: the Scientific Committee on Consumer Safety (SCCS) and the Scientific Committee on Health, Environmental and Emerging Risks (SCHEER) and are made up of scientists appointed in their personal capacity.

In addition, the Commission relies upon the work of the European Food Safety Authority (EFSA), the European Medicines Agency (EMA), the European Centre for Disease prevention and Control (ECDC) and the European Chemicals Agency (ECHA).

#### **SCCS**

The Committee shall provide Opinions on questions concerning all types of health and safety risks (notably chemical, biological, mechanical and other physical risks) of non-food consumer products (for example: cosmetic products and their ingredients, toys, textiles, clothing, personal care and household products such as detergents, etc.) and services (for example: tattooing, artificial sun tanning, etc.).

#### Scientific Committee members

Bernauer Ulrike, Bodin Laurent, Chaudhry Mohammad Qasim, Coenraads Pieter-Jan, Dusinska Maria, Ezendam Janine, Gaffet Eric, Galli Corrado Lodovico, Granum Berit, Panteri Eirini, Rogiers Vera, Rousselle Christophe, Stępnik Maciej, Vanhaecke Tamara, Wijnhoven Susan

### Contact

Doi:

European Commission
Health and Food Safety
Directorate C: Public Health, Country Knowledge, Crisis Management
Unit C2 – Country Knowledge and Scientific Committees
L-2920 Luxembourg
SANTE-C2-SCCS@ec.europa.eu

C	European	Union,	2018	

ISSN ISBN

The Opinions of the Scientific Committees present the views of the independent scientists who are members of the committees. They do not necessarily reflect the views of the European Commission. The Opinions are published by the European Commission in their original language only.

ND-

http://ec.europa.eu/health/scientific\_committees/consumer\_safety/index\_en.htm

# **TABLE OF CONTENTS**

1.	MANDATE FROM THE EUROPEAN COMMISSION	5
2.	OPINION	7
3.	CONCLUSION	7
4.	MINORITY OPINION	8
5.	REFERENCES	9
6.	GLOSSARY OF TERMS	36
7	LIST OF ABBREVIATIONS	36

# 1. MANDATE FROM THE EUROPEAN COMMISSION

# **Background**

Biphenyl-2-ol and its salts, covering o-Phenylphenol (OPP), Sodium o-phenylphenate, Potassium o-phenylphenate, MEA o-phenylphenate (CAS n. 90-43-7, 132-27-4, 13707-65-8, 84145-04-0) as preservatives are regulated in Annex V, entry 7 of the Cosmetics Regulation (EC) n. 1223/2009 at a maximum concentration of 0.2 % (as phenol).

The substance MEA o-phenylphenate is also regulated through entry 61 of Annex III on Monoalkylamines, monoalkanolamines and their salts which are restricted for use in cosmetic products to a maximum secondary amine content of 0,5% in ready for use preparation.

The SCCS Committee adopted an opinion on o-Phenylphenol, Sodium o-phenylphenate and Potassium o-phenylphenate (SCCS/1555/15) in June 2015, later revised on 15 December 2015, with the following conclusion:

"o-Phenylphenol as preservative with a maximum concentration of 0.2 % in leave-on cosmetic products is not safe. Also, in view of further exposures including non-cosmetic uses, the maximum concentration of o-Phenylphenol in leave-on cosmetic products should be lowered.

However, the proposed maximum use concentration of up to 0.15% by the applicant can be considered safe.

The use of o-Phenylphenol as preservative with a maximum concentration of 0.2 % in rinse-off cosmetic products is considered safe.

Based on the information provided, no conclusions of safe use can be drawn for Sodium ophenylphenate and Potassium o-phenylphenate."

Following this SCCS opinion, several Member States raised specific concerns as regards the potential risk to human health of Sodium o-phenylphenate, Potassium o-phenylphenate and MEA o-phenylphenate.

In December 2017, the German Federal Institute for Risk Assessment submitted a document providing information on the safety of Sodium o-phenylphenate, Potassium o-phenylphenate and MEA o-phenylphenate when compared to OPP.

# **Terms of reference**

- 1. Does SCCS consider Sodium o-phenylphenate, Potassium o-phenylphenate and MEA o-Phenylphenate safe at the current use as preservatives with a maximum concentration of 0.2 % (as phenol)?
- 2. Does the SCCS consider that the same conclusion for OPP, as reported in SCCS/1555/15, may also be applied to Sodium o-phenylphenate, Potassium o-phenylphenate and MEA o-Phenylphenate concerning the proposed maximum use concentration (i.e. 0.15% in leave-on and 0.2% in rinse-off cosmetic products as preservatives)?

# 2. OPINION

The SCCS opinion (SCCS/1555/15) relates to the safety of o-Phenylphenol (OPP), sodium-OPP, and potassium-OPP for use as preservatives, and is based on the information received in response to the Commission's call for further data. The available information mainly relates to OPP and sodium-OPP, limited information on potassium-OPP, and virtually no information on MEA-OPP. Also the document provided by the German Federal Institute for Risk Assessment did not contain any further information on sodium-OPP, potassium-OPP or MEA-OPP.

#### 3. CONCLUSION

1. Does SCCS consider Sodium o-phenylphenate, Potassium o-phenylphenate and MEA o-Phenylphenate safe at the current use as preservatives with a maximum concentration of 0.2 % (as phenol)?

Due to the lack of relevant information, the SCCS is unable to answer the question on the safe use level of sodium-OPP, potassium-OPP and MEA-OPP. In SCCS view, a direct comparison between the safety of o-phenylphenate (OPP) and its 3 compounds cannot be made due to the following concerns:

- From the limited available information, it is clear that both sodium and potassium salts
  of OPP have much higher water solubility than OPP (no information available on MEAOPP). This can potentially alter their absorption and biokinetics, compared to OPP.
- From the available information, the SCCS has noted that, compared to the strongly skin irritating nature of OPP, both sodium OPP and potassium-OPP are corrosive to the skin, and sodium OPP is also corrosive to the eye. This indicates that both sodium and potassium salts of OPP may have greater skin penetration and potentially more potent toxic effects than OPP due to higher systemic exposure. No relevant information on skin irritation is available for MEA-OPP but the presence of monoethyleneamine (MEA) moiety can also be expected to alter the skin absorption and biokinetics of MEA-OPP compared to OPP, and as a consequence also systemic exposure. For OPP, the SCCS has derived dermal absorption value of 45% from toxicokinetic information. However, such information is not available for sodium-OPP, potassium-OPP or MEA-OPP, and data would be needed to allow drawing any comparisons from the safe use levels of OPP.
- The available in vivo studies in rat have also indicated that the adverse effects of OPP and its sodium salt are different. For the sodium salt, there is clear indication that the substance is more potent with respect to urinary bladder carcinoma and data point to mechanistic differences between OPP and SOPP. Amongst other factors, SOPP leads to higher sodium concentrations in urine and also to higher urinary pH. There is insufficient dose-response data available to draw a conclusion on the possibility of setting a threshold for sodium-OPP induced toxicity. The currently available data are also not sufficient to exclude such a possibility for the other two compounds (potassium-OPP and MEA-OPP). Long-term repeat dose studies have pointed out to a threshold between 35 and 40 mg/kg bw/d for OPP, but due to the lack of dose-response data, a threshold for sodium-OPP, potassium-OPP or MEA-OPP cannot be derived.

2. Does the SCCS consider that the same conclusion for OPP, as reported in SCCS/1555/15, may also be applied to Sodium o-phenylphenate, Potassium o-phenylphenate and MEA o-Phenylphenate concerning the proposed maximum use concentration (i.e. 0.15% in leave-on and 0.2% in rinse-off cosmetic products as preservatives)?

For the reasons given above, the SCCS considers that the same conclusions on the safe use levels of OPP cannot be applied as such to sodium-OPP, potassium-OPP or MEA-OPP for use in rinse-off and leave-on cosmetic products.

Based on the available information, the SCCS is of the opinion that a potential risk to human health from the use of sodium-OPP and potassium-OPP as preservatives in cosmetic products cannot be excluded.

Although the safety of MEA-OPP was not evaluated in the Opinion SCCS/1555/15, the SCCS has a similar view that a potential risk from its use as preservative in cosmetic products cannot be excluded in the absence of relevant data.

# 4. MINORITY OPINION

/

\_\_\_\_\_\_

#### 5. REFERENCES

#### From Opinion SCCS/1555/15

- 1. Adams, R.M. Allergic Contact Dermatitis due to O-Phenylphenol. Contact Dermatitis 7, 332, 1981
- Andersen, K.E., Hamann, K. The Sensitizing Potential of Metalworking Fluid Biocides (Phenolic and Thiazolic Compounds) in The Guinea-Pig Maximization Test in Relation to Patch-Test Reactivity In Eczema Patients. Food Chem. Toxicol. 22, 655-660, 1984
- 3. Appel, K.E. The Carcinogenicity of the Biocide Ortho-Phenylphenol. Arch Toxicol 74, 61-71, 2000
- 4. Bajaj, K.L., Miller, I.R., Bhatia, I.S. Metabolism of 2-Hydroxybiphenyl & 4-Hydroxybiphenyl in Albino Mice. Indian J. Exp. Biol. 14, 329-331, 1976
- Balakrishnan & Eastmond, Micronuclei and cell proliferation as early biological markers of ortho-phenylphenol-induced changes in the bladder of male F344 rats.
   Food Chemical Toxicology 44, 1340-1347, 2006
- Balakrishnan, S., Uppala, P.T., Rupa, D.S., Hasegawa, L., Eastmond, D.A. Detection of Micronuclei, Cell Proliferation and Hyperploidy in Bladder Epithelial Cells Of Rats Treated With O-Phenylphenol. Mutagenesis 17, 89-93, 2002
- 7. Bartels, M.J. Mcnett, D.A. Quantitation of Orthophenylphenol Metabolites in Rat Urine Samples from A <sup>32</sup>P-Postlabeling Study. Dow Chemical, Report No. K-001024-062, 1996
- 8. Bartels, M.J., Brzak, K.A., Mcnett, D.A., Shabrang, S.N. Ortho-Phenylphenol (OPP): Limited Metabolism Study in Human. Dow Chemical, Report No. K-0011024-059, 1997
- 9. Bartels, M.J., Mcnett, D.A., Timchalk, C., Mendrala, A.L., Christenson, W.R., Sangha, G.K., Brzak, K.A., Shabrang, S.N. Comparative Metabolism of Ortho-Phenylphenol in Mouse, Rat and Man. Xenobiotica 28, 579-594, 1998
- 10. Berdasco, N.M. Orthophenylphenol: Dermal Sensitization Potential in the Hartley Albino Guinea Pig. Dow Chemical, Report No. K-001204-048, 1991
- 11. Blair, R.M., Fang, H., Branham, W.S., Hass, B.S., Dial, S.L., Moland, C.L., Tong, W., Shi, L., Perkins, R., Sheehan, D.M. The Estrogen Receptor Relative Binding Activities of 188 Natural And Xenochemicals: Structural Diversity Of Ligands. Toxicol. Sci. 54, 138-153, 2000
- 12. Bomhard EM et al. O-Phenylphenol and its Sodium and Potassium Salts: A toxicological Assessment. Critical Reviews in toxicology, 32(6):551-626(2002)

- 13. Bomhard et al., o-Phenyolphenol and its sodium and potassium salts: a toxicological assessment. Critical Reviews in Toxicology 32, 551-626, 2002
- 14. Bomhard, E. O-Phenylphenol Kaliumsalz Untersuchungen zur Akuten Oralen Toxizität an Männlichen und Weiblichen Wistar-Ratten. Bayer Ag, Report No. 17260, 1988
- 15. Bomhard, E. O-Phenylphenol Kaliumsalz Untersuchungen zur Akuten Dermalen Toxizität An Männlichen und Weiblichen Wistar-Ratten. Bayer Ag, Report No. 20817, 1991b
- 16. Bomhard, E. Preventol O Extra (Schuppen) Untersuchungen zur Akuten Dermalen Toxizität an Männlichen und Weiblichen Wistar-Ratten. Bayer Ag, Report No. 19831, 1991a
- 17. Boutwell, R.K., Bosch, D.K. The Tumor-Promoting Action of Phenol and Related Compounds for Mouse Skin. Cancer Res. 19, 413-424, 1959
- 18. Brasch, J., Henseler, T., Frosch, P. Patch Test Reactions to a Preliminary Preservative Series. Dermatosen, 41, 71-76, 1993
- 19. Brendler, S.Y. Preventol O-Extra: Mutagenicity Study for the Detection of Induced Forward Mutations in the Cho-HGPRT Assay In Vitro. Bayer Ag, Report No. 21278, 1992
- 20. Brendler-Schwaab, S.Y. Preventol O-Extra: Comet Assay in Vivo in Mouse Liver and Kidney. Bayer Ag, Report No. 30130, 2000
- 21. Brusick: Analysis of genotoxicity and the carcinogenic mode of action for orthophenylphenol. Environmental Molecular Mutagenesis 45, 460-481, 2005
- 22. Burke, M.D., Bridges, J.W. Biphenyl Hydroxylations and Spectrally Apparent Interactions with Liver Microsomes from Hamsters Pre-Treated With Phenobarbitone And 3-Methylcholanthrene. Xenobiotica 5, 357-376, 1975
- 23. Carreon, R.E., New, M.A. Dowicide 1: Acute Percutaneous Absorption Potential. Dow Chemical, Report No. K-1024-(37), 1981
- 24. Christenson, W.R., Wahle, B.S., Bartels, M.J., Cohen, S.M. Technical Grade Ortho-Phenylphenol: A 32-P-Postlabeling Study to Examine the Potential for the Formation of DNA Adducts in the Urinary Bladder of the Male Rat. Bayer Corp., Report No. 94-972-Av, 1996b
- 25. Christenson, W.R., Wahle, B.S., Cohen, S.M. Technical Grade Ortho-Phenylphenol: A Special Subchronic Dietary Study to Examine the Mechanism of Urinary Bladder Carcinogenesis in the Male Rat. Bayer Corp., Report No. 92-972-Ms, 1996a
- 26. Cline, J.C., Mcmahon, R.E. Detection of Chemical Mutagens. Use of Concentration Gradient Plates in a High Capacity Screen. Res. Comm. Chem. Pathol. Pharmacol. 16, 523-533, 1977

- 27. Cnubben, N. Comparative in vitro in vivo Percutaneos Penetration of Fungicide Ortho-Phenylphenol Regulatory Toxicology and Pharmacology, 35, 198-208, 2002
- 28. Cohen, S.M. Role of Urinary Physiology and Chemistry in Bladder Carcinogenesis. Food Chem. Toxicol. 33, 715-730, 1995
- 29. Cohen, S.M. Urinary Bladder Carcinogenesis. Toxicol. Pathol. 26, 121-127, 1998
- 30. Cohen, S.M., Cano, M., Anderson, T., Garland, E.M. Extensive Handling of Rats Leads to Mild Urinary Bladder Hyperplasia. Toxicol. Pathol. 24, 251-257, 1996
- 31. Cohen, S.M., Cano, M., Earl, R.A., Carson, S.D., Garland, E.M. A Proposed Role for Silicates and Protein in the Proliferative Effects of Saccharin on the Male Rat Urothelium. Carcinogenesis 12, 1551-1555, 1991
- 32. Cosse, P.F., Stebbins, K.E., Stott, W.T., Johnson, K.A., Atkin, L. Ortho-Phenylphenol: Palatability/ Probe, Four-Week and one-year Oral Toxicity Studies in Beagle Dogs. Dow Chemical, Report No. K-001024-039, 1990
- 33. Eastmond, D.A. Induction of Micronuclei and Aneuploidy By The Quinone-Forming Agents Benzene And Ophenylphenol. Toxicol. Letters 67, 105-118, 1993
- 34. Eastmond, D.A. Personal Communication March 2002
- 35. EFSA: Peer review of the pesticide risk assessment of the active substance 2-phenylphenol. EFSA Scientific Report 217, 1-67, 2008
- 36. Eigenberg, D.A. Two-Generation Dietary Reproduction Study in Rats Using Orthophenylphenol. Mobay Corp., Report No. 85-671-02; 1990
- 37. Eigenberg, D.A., Lake, S.A. A Two-Generation Dietary Reproduction Study in Sprague-Dawley Rats Using Technical Grade Orthophenylphenol. Bayer Corp., File No. 7788; 1995
- 38. Eigenberg, D.A., Lake, S.G., Sangha, G.K., Thyssen, J.H. Evaluation Of The Reproductive Toxicity Of O-Phenylphenol (OPP) in a Two-Generation Rat Reproductive Toxicity Study. Fundam. Appl. Toxicol. 36, 356, 1997 (Abstract)
- 39. EPA Environmental Protection Agency Notice: Endocrine Disruptor Screening Program [Oppts-42206; Frl-6021-3]. Federal Register: August 11, 1998 [63, No. 154, 42852-42855]
- 40. EPA Environmental Protection Agency Special Report on Environmental Endocrine Disruption: An Effects Assessment and Analysis. Washington D.C., 1997
- 41. Ernst, W. Umwandlung und Ausscheidung von 2-Hydroxydiphenyl Bei der Ratte. Arzneim. Forsch. 15, 632-636, 1965
- 42.EU European Commission DG ENV. Towards the Establishment of a Priority List of Substances for Further Evaluation of Their Role in Endocrine Disruption. June 21, 2000

- 43. EU MRC Institute for Environment and Health European. Workshop on the Impact of Endocrine Disrupters on Human Health and Wildlife. Reports of Proceedings Weybrigde, U.K.; EUR 17549, 1996
- 44. FAO/WHO Plant Production and Protection Paper, No. 102, 49, 1990
- 45. FAO/WHO Plant Production And Protection Paper, No. 153, 169, 1999
- 46. FAO/WHO Plant Production and Protection Paper, No. 68, 37, 1985
- 47. Freyberger, A. O-Phenylphenol Interactions Of O-Phenylphenol (OPP) and its Metabolites with Microsomal Prostaglandin-H-Synthase: Possible Implications for OPP-Induced Tumor Formation in The Rat Urinary Bladder. Bayer Ag, Report No. 22788, 1994
- 48. Freyberger, A., Degen, S.A. Inhibition of Prostaglandin-H-Synthase by O-Phenylphenol and its Metabolites Arch. Toxicol. 72, 637-644, 1998
- 49. Fudan University; Department of Toxicology, School of Public Health, Study Report August 1, 2006
- 50. Fujii, T., Hiraga, K. Carcinogenicity Testing of Sodium Orthophenylphenate in F344 Rats. J. Saitana Med. School 12, 277-287, 1985
- 51. Fujii, T., Mikuriya, H., Hayashida, S., Hiraga, K. Enhancing Effects of Thiabendazole on Urinary Bladder Carcinogenesis in Rats Fed Diet Containing Low-Dose of Sodium O-Phenylphenate for 13 Week. Ann. Rep. Tokyo. Metr. Res. Lab. P.H. 37, 411-414, 1986b
- 52. Fujii, T., Mikuriya, H., Kamiya, N., Hiraga, K. Enhancing Effect of Thiabendazole on Urinary Bladder Carcinogenesis Induced by Sodium Ophenylphenate in F344 Rats. Food Chem. Toxic. 24, 207-211, 1986a
- 53. Fujii, T., Mikuriya, H., Sasaki, M. Chronic Oral Toxicity and Carcinogenicity Study of Thiobendazole in Rats. Food Chem. Toxicol. 29, 771-775, 1991
- 54. Fujii, T., Mikuriya, H., Yano, N., Yuzawa, K:, Nagasawa, A., Fukumori, N., Tada, Y., Sasaki, M. Subacute Toxicity Test of Ceramics by Gavage to Rats and Influence of Ceramics on Urinary Bladder Carcinogenesis Induced by O-Phenylphenol. Ann. Rep. Tokyo Metr. Res. Lab. P.H. 41, 233-244, 1990
- 55. Fujii, T., Mikuriya, H., Yoneyama, M., Yano, N., Yuzawa, K., Nagasawa, A., Sasaki, M. Dietary Toxicity Test of O-Phenylphenol and Sodium Hydrogencarbonate by Simultaneous Administration to Mice for 52 Weeks. Ann. Rep. Tokyo. Metr. Res. Lab. P.H. 40, 298-306, 1989a
- 56. Fujii, T., Mikuriya, H., Yoneyama, M., Yano, N., Yuzawa, K., Nagasawa, A., Sasaki, M. Dietary Toxicity of O-Phenylphenol, Thiabendazole, and Sodium Hydrogencarbonate by Simultaneous Administration to Mice for 52 Weeks. Ann. Rep. Tokyo. Metr. Res. Lab. P.H. 40, 307-315, 1989b

- 57. Fujii, T., Nakamura, K., Hiraga, K. Effects of pH on the Carcinogenicity of O-Phenylphenol and Sodium O-Phenylphenate in the Rat Urinary Bladder. Food Chem. Toxic. 25, 359-362, 1987
- 58. Fujita, H., Kojima, A., Sasaki, M., Hiraga, K. Assay of Host-Mediated Mutagenicity by S. Typhimurium and E. Coli in Rats Fed Sodium O-Phenylphenate (OPP-Na). Ann. Rep. Tokyo Metr. Res. Lab. P.H. 35, 431-435, 1984
- 59. Fujita, H., Kojima, A., Sasaki, M., Hiraga, K. Mutagenicity Test of Antioxidants and Fungicides with Salmonella Typhimurium Ta97a And Ta102. Ann. Rep. Tokyo Metr. Res. Lab. P.H. 36, 413-417, 1985
- 60. Fukumori, N., Sasaki, M. Comparison of Ultrastructural Changes between O-Phenylphenol (Opp) Kidney Damage and Sucrose-Nephrosis in the Rat Proximal Tubules. Ann. Rep. Tokyo Metr. Res. Lab. P.H. 37, 399-406, 1986
- 61. Fukumori, N., Sasaki, M. Ultrastructural Alterations of the Proximal Tubules Induced by O-Phenylphenol (OPP) in the Rat Kidney. Ann. Rep. Tokyo Metr. Res. Lab. P.H. 36, 400-408, 1985
- 62. Fukumori, N., Sasaki, M., Hiraga, K. Effects of Sodium O-Phenylphenate (OPP-Na) on Ultra-Structure of Bladder Epithelium in Rats. Ann. Rep. Tokyo Metr. Res. Lab. P.H. 35, 416-424, 1984
- 63. Fukumori, N., Sasaki, M., Hiraga, K. Ultrastructural Alterations of Hepatocytes in Male Rats Fed With Sodium O-Phenylphenate (Oppna). Ann. Rep. Tokyo Metr. Res. Lab. P.H. 34, 329-336, 1983
- 64. Fukushima, S., Hasegawa, R., Kurata, Y., Okuda, M., Hatano, A., Ito, N. Histopathological and Ultrastructural Analysis of Urinary Bladder Lesions in Animals Induced by Sodium O-Phenylphenate. Proc. Jap. Cancer Ass., 41 Ann. Mtg., 314, 1982 (Abstract)
- 65. Fukushima, S., Inoue, T., Uwagawa, S., Shibata, M.A., Ito, N. Co-Carcinogenic Effects of NaHCO3 on O-Phenylphenol-Induced Rat Bladder Carcinogenesis. Carcinogenesis 10, 1635-1640, 1989
- 66. Fukushima, S., Kurata, Y., Ogiso, T., Okuda, M., Miyata, Y., Ito, N. Pathological Analysis of the Carcinogenicity of Sodium O-Phenylphenate and O-Phenylphenol. Oncology 42, 304-311, 1985
- 67. Fukushima, S., Kurata, Y., Shibata, M.A., Ikawa, E., Ito, N. Promoting Effect of Sodium O-Phenylphenate and O-Phenyl-Phenol on Two-Stage Urinary Bladder Carcinogenesis In Rats. Jpn. J. Cancer Res.(Gann) 74, 625-632, 1983
- 68. Fukushima, S., Shibata, M.A., Kurata, Y., Tamano, S., Masui, T. Changes in the Urine and Scanning Electron Microscopically Observed Appearance of the Rat

- Bladder Following Treatment With Tumor Promotors. Jpn. J. Cancer Res.(Gann) 77, 1074-1082, 1986
- 69. Funahashi, M., Arai, M., Takahashi, H., Hibino, T., Morphological Characteristics of Rabbit Urinary Bladder Tumors Induced by BBN and OPP-Na. Fujita Gakuen Hoken Eisei Daigaku, Toyoaki 11, 187-189, 1987
- 70. Gaches, C. G. Woodworm and The Bladder. Proc. R. Soc. Med. 68, 525-527, 1975
- 71. Gbodi, T.A., Oehme, F.W. The Fate Of Phenol, O-Phenylphenol and Disophenol in Rats. Toxicol. Appl. Pharmacol. 45, 223, 1978 (Abstract)
- 72. Geier, J., Kleinhans, D., Peters, K.-P. Kontaktallergien durch Industriell Verwendete Biozide. Dermatosen / Occup. Environ. 44, 154-159, 1996
- 73. Geter et al. 2009a: Profiling Ortho-Phenylphenol-Induced Molecular, Cellular, and Biochemical Changes in B6C3F1 Male Mice. Toxicology & Environmental Research and Consulting (TERC) Study ID: 080538; The Dow Chemical Company. 9 October 2009
- 74. Geter et al. 2009b: Ortho-Phenylphenol (OPP): An assessment of the relevance of hepatocellular tumors in mice to humans. Evaluation dated 29 October 2009
- 75. Gilbert, K.S. Dowicidetm1 Antimicrobial; Dermal Sensitization Potential In The Hartley Albino Guinea Pig. Dow Chemical, Report No. K-001024-057e, 1994b
- 76. Gilbert, K.S. Dowicidetm1 Antimicrobial; Primary Dermal Irritation Study in New Zealand White Rabbits. Dow Chemical, Report No. K-001024-057b, 1994a
- 77. Gilbert, K.S., Crissman, J.W. Dowicidetm1 Antimicrobial; Acute Oral Toxicity Study in Fischer 344 Rats. Dow Chemical, Report No. K-001024-057a, 1994
- 78. Gilbert, K.S., Stebbins, K.E. Dowicide A Antimicrobial: Acute Oral Toxicity Study In Fischer 344 Rats. Dow Chemical, Report No. K-001025-014a, 1994
- 79. Gilbert; K.S. Dowicide A Antimicrobial: Dermal Sensitization Potential in Hartley Albino Guinea Pigs. Dow Chemical, Report No. K-0011025-014e, 1994c
- 80. Goh, C.L., Yuen, R. A Study of Occupational Skin Disease in the Metal Industry (1986-1990). Ann. Acad. Med. Singapore 23, 639-644, 1994
- 81. Grattan, C.E.H., English, J.S.C., Foulds, I.S., Rycroft, R.J.G. Cutting Fluid Dermatitis. Contact Dermatitis, 20, 372-376, 1989
- 82. Grether, T., Brunn, H., Laib, R.J. 32P-Postlabelling Method as a Sensitive Indicator for Analysis of Genotoxicity of Biphenyl Derivatives. Arch. Toxicol. 63, 423-424, 1989a
- 83. Grether, T., Rogiers, V., Laib, R.J. Analysis of Genotoxicity of Hydroxylated and Chlorinated Biphenyl Derivatives in vitro and in vivo. Naunyn-Schmiedeberg's Arch. Pharmacol. Suppl. To Vol. 339, R 23, 1989b (Abstract)

- 84. Groot, A.P. De, Feron, V.J., Immel, H., Rats R. Induction of Hyperplasia in the Bladder Epithelium of Rats by a Dietary Excess of Acid or Base: Implications for Toxicity/Carcinogenicity Testing. Food Chem. Toxic. 26, 425-434, 1988
- 85. Gucklhorn, I.R. Antimicrobials in Cosmetics. Manufacturing Chemist Aerosol News 40, 23-30, 1969
- 86. Hagedorn-Leweke, U., Lippold, B.C. Absorption of Sunscreens and Other Compounds through Human Skin in vivo: Derivation of a Method to Predict Maximum Fluxes. Pharm. Res. 12, 1354-1360, 1995
- 87. Hagiwara, A., Shibata, M., Hirose, M., Fukushima, S., Ito, N. Long-Term Toxicity and Carcinogenicity Study of Sodium O-Phenylphenate in B6C3F1 Mice. Food Chem. Toxic. 22, 809-814, 1984
- 88. Halpaap-Wood, K., Horning, E.C., Horning, M.G. The Effect of 3-Methylcholanthrene, Aroclor 1254, and Phenobarbital Induction on the Metabolism of Biphenyl by Rat and Mouse 9000g Supernatant Liver Fractions. Drug Metab. Disp. 9, 103-107, 1981
- 89. Hanada, S. Studies on Food Additives Diphenyl and O-Phenylphenol From Standpoint of Public Health. Toxicological Studies of Diphenyl and O-Phenylphenol. J. Nagoya City Univ. Med. Ass. 28, 983-995,1977
- 90. Harbell, J.W. O-Phenylphenol Mouse Lymphoma Assay (L5178y Tk +/-). Microbiological Associates Inc., (Lab. Study Number MI-Nci #246);1989a
- 91. Harbell, J.W. O-Phenylphenol, Sodium Salt Tetrahydrate Mouse Lymphoma Assay (L5178y Tk +/-). Microbiological Associates Inc., (Lab. Study Number MI-Nci #247); 1989b
- 92. Harke, H.-P., Klein, H., Zur Frage der Resorption von 2-Phenylphenol aus Waschenden Haende-Desinfektionsmitteln. Zbl. Bakt. Hyg., I. Abt. Orig. B 174, 274-278, 1981
- 93. Hasegawa, R., Cohen, S.M. The Effect of Different Salts of Saccharin on the Rat Urinary Bladder. Cancer Lett. 30, 261-268, 1986
- 94. Hasegawa, R., Fukuoka, M., Takahashi, T., Yamamoto, A., Yamaguchi, S., Shibata, M.A., Tanaka, A., Fukushima, S. Sex Differences in O-Phenylphenol and Sodium O-Phenylphenate Rat Urinary Bladder Carcinogenesis: Urinary Metabolites and Electrolytes Under Conditions of Aciduria and Alkalinuria. Jpn. J. Cancer Res. 82, 657-664, 1991
- 95. Hasegawa, R., Furukawa, F., Toyoda, K., Sato, H., Shimoji, N., Takahashi, M., Hayashi, Y. In situ Freezing of the Urinary Bladder: A Trigger Of Rapid Development Of Sodium-O-Phenylphenate- Induced Urinary Bladder Tumors In The Rat. Carcinogenesis 10, 571-575, 1989a

- 96. Hasegawa, R., Furukawa, F., Toyoda, K., Sato, H., Takahashi, M., Hayashi, Y. Urothelial Damage and Tumor Initiation by Urinary Metabolites of Sodium-O-Phenylphenate in the Urinary Bladder of Female Rats. Jpn. J. Cancer Res. 81, 483-488, 1990a
- 97. Hasegawa, R., Kaji, N., Furukawa, F., Fukuoka, M., Kuzunishi, K., Takahashi, M., Hayashi, Y. Investigation of Bladder Cancer Initiation Effects of OPP-Na Metabolites in Rat Urine. Proc. J. Cancer Assoc., 91, 1988, 235 (Abstract)
- 98. Hasegawa, R., Nakaji, Y., Kurokawa, Y., Tobe, M. Acute Toxicity Tests On 113 Environmental Chemicals. Sci. Rep. Inst. Tohuku Univ. 36, 1-4, 1989b
- 99. Hasegawa, R., Takahashi, S., Asamoto, M., Shirai, T., Fukushima, S. Species Differences In Sodium O-Phenylphenate Induction of Urinary Bladder Lesions. Cancer Lett. 50, 87-91, 1990b
- 100. Haworth, S., Lawlor, T., Mortelmans, K., Speck, W., Zeiger, E. Salmonella Mutagenicity Test Results for 250 Chemicals. Environ. Mut. Suppl.1, 3, 1983
- 101. Henschke, P., Almstadt, E., Lüttgert, S., Appel, K.E. Metabolites of the Biocide O-Phenylphenol Generate Oxidative DNA Lesions in V79 Cells. Arch. Toxicol. 73, 607-610, 2000
- 102. Hiraga, K, Fujii, T. Induction of Tumours of the Urinary Bladder in F344 Rats by Dietary Administration of O-Phenylphenol. Food Chem. Toxicol. 22, 865-870, 1984
- 103. Hiraga, K., Fujii, T. Induction of Tumours of the Urinary System in F344 Rats by Dietary Administration of Sodium O-Phenylphenate. Food Cosmet. Toxicol. 19, 303-310, 1981
- 104. Hirayama, T., Nohara, M., Shindo, H., Fukui, S. Mutagenicity Assays of Photochemical Reaction Products Of Biphenyl (BP) and O-Phenylphenol (OPP) With Nox. Chemosphere 10, 223-228, 1981
- 105. Hodge et al., Toxicological studies of orthophenylphenol (Dowcide 1). Journal of Pharmacology and experim. Therapeut. 104, 202-210, 1952
- 106. Hodge, H.C., Maynard, E.A., Blanchet, H.J., Spencer, H.C., Rowe, V.K. Toxicological Studies of Orthophenylphenol (Dowicide 1). J. Pharmacol. Exp. Ther. 104, 202-210, 1952
- 107. Honma, Y., Kakizoe, T., Komatsu, H., Niijima, T., Sugimura, T. Increased Agglutinability of Bladder Epithelial Cells by Concanavalin A in Rats Fed Several Biphenyl Derivatives. J. Cancer Res. Clin. Oncol. 106, 176-178, 1983
- 108. Horvath, E., Levay, G., Pongracz, K., Bodell, W.J. Peroxidative Activation of Ortho-Phenylhydroquinone Leads to the Formation of DNA Adducts in HI-60 Cells. Carcinogenesis 13, 1937-1939, 1992

- 109. IARC: Ortho-Phenylphenol and its Sodium Salt. Monographs on the Evaluation of Carcinogenic Risks of Chemicals to Humans, Miscellaneous Pesticides 30, 329-344, 1983
- 110. IARC: Ortho-Phenylphenol and its Sodium Salt. Monographs on the Evaluation of Carcinogenic Risks To Humans 73, 451-480, 1999
- 111. IARC: Sex Hormones. Monographs on the evaluation of Carcinogenic Risks of Chemicals to Humans. 21, 1979
- 112. Iguchi, S., Takahashi H., Fugii, T., Fukumori, N., Mikuriya, H., Tada, Y., Yuzawa, K., Hiraga, K. Subchronic Toxicity of O-Phenylphenol (Opp) by Food Administration to Rats. Ann. Rep. Tokyo Metr. Res. Lab. P.H. 35, 407-415, 1984
- 113. Iguchi, S., Tayama, K., Hiraga, K. Subacute Toxicity of Sodium O-Phenylphenate by Food Administration to Rats. Ann. Rep. Tokyo Metr. Res. Lab. P.H. 30, 67-79, 1979
- 114. Innes, J.R.M., Ulland, B.M., Valerio, M.G., Petrucelli, L., Fishbein, L., Hart, E.R., Palotta, A.J., Bates, R.R., Falk, H.L., Gart, J.J., Klein, M., Mitchell, I., Peters, J. Bioassay of Pesticides and Industrial Chemicals for Tumorigenicity in Mice: A Preliminary Note. J. Natl. Cancer Inst. 42, 1101-1114, 1969
- 115. Inoue, S., Yamamoto, K., Kawanishi, S. DNA Damage Induced by Metabolites of O-Phenylphenol in the Presence of Copper(II) ons. Chem. Res. Toxicol. 3, 144-149, 1990
- 116. Inoue, T. Pathological Studies on Urinary Bladder Carcinogenesis of O-Phenylphenol and its Sodium Salt in Male Rats. J. Nagoya City Univ. Med. Assoc. 44, 463-478, 1993
- 117. Inoue, T., Shibata, M., Uwa, K., Asamoto, S., Osaki, T., Fukushima, A. Synergistic Effect Of NaHCO3 on Rat Bladder Carcinogenesis by OPP. Proc. J. Cancer Assoc., 91, 1988 (Abstract)
- 118. Inui, N., Nishi, Y., Iwata, K. Dose-Response Relationships for Mutations Induced in Embryo Cells after Treatment of Pregnant Hamsters. In: Problems of Threshold in Chemical Mutagenesis. Eds.: Tazima, Y. Et Al., 35-39, 1984
- 119. Ishidate, M. (Ed.) Chromosomal Aberration Test in Vitro. L.I.C., Inc., Tokyo, 1987
- 120. Ishidate, M., Harnois, M.C., Sofuni, T. A Comparative Analysis of Data on the Clastogenicity of 951 Chemical Substances Tested in Mammalian Cell Cultures. Mutat. Res. 195, 151-213, 1988
- 121. Ishidate, M., Sofuni, T., Yoshikawa, K., Hayashi, M., Nohmi, T., Sawada, M., Matsuoka, A. Primary Mutagenicity Screening Of Food Additives Currently Used In Japan. Food Chem. Toxic. 22, 623-636, 1984

- 122. Ishidate, M., Yoshikawa, K., Sofuni, T., Mutagenicity Tests on OPP, OPP-Sodium Salt, and Their Possible Metabolites. Unpublished Report From the Division of Mutagenesis, Biological Safety Research Center, National Institute of Hygienic Sciences, Tokyo, Japan, 1983 Submitted To Who By Dow Chemical Co., Midland, Mt., USA Cited according to: FAO/WHO Evaluations 1985, Toxicology
- 123. Ito, K., Nishitani, K., Hara, I. A Study of Cases of Leucomelanodermatosis due to Phenyl-Phenol Compounds. Bull. Pharm. Res. Institute 76, 5-13, 1968
- 124. Ito, N., Fukushima, S., Shirai, T., Hagiwara, A., Imaida, K. Drugs, Food Additives and Natural Products as Promoters in Rat Urinary Bladder Carcinogenesis. Iarc Sci. Publ. 56, 399-407, 1984
- 125. Ito, N., Hasegawa, R., Imaida, K., Takahashi, S., Shirai, T. Medium-Term Rat Liver Bioassay for Rapid Detection of Carcinogens and Modifiers of Hepatocarcinogenesis. Drug Metab. Rev. 26, 431-442, 1994
- 126. Itoh, S., Ueda, H., Naasaka, T., Sumitomo, H. Evaluating Variation Of Estrogenic Effect by Drinking Water Chlorination With The MvIn Assay. Water Sci. Technol. 42, 61-69, 2000
- 127. JECFA (Joint FAO/WHO Expert Committee on Food Additives) fifty-fifth meeting, Geneva, 6-15 June 2000
- 128. JMPR, Pesticide Residues In Food 1985 Report On The Joint Meeting of the FAO Panel of Experts on Pesticide Residues in Food and the Environment and a WHO Expert Group on Pesticide Residues. Geneva, 37, (1985)
- 129. John, J.A., Murray, F.J., Rao, K.S., Schwetz, B.A. Teratological Evaluation of Orthophenylphenol in Rats. Fundam. Appl. Toxicol. 1, 282-285, 1981
- 130. Kabashima, J., Ichikawa, H., Nakao, T. Effect of Sodium O-Phenylphenol (OPP-Na) on Acid Phosphatase in Rat Urine. Ann. Rep. Tokyo Metr. Res. Lab. P.H. 34, 309-311, 1983
- 131. Kabashima, J., Nakao, T. Effect of Sodium O-Phenylphenate (OPP-Na) on Acid Phosphatase in Rat Urine. Ann. Rep. Tokyo Metr. Res. Lab. P. H. 32, 61-64, 1981
- 132. Kahn, G. Depigmentation Caused by Phenolic Detergent Germicides. Arch. Dermatol. 102, 177-187, 1970
- 133. Kaneda, M., Teramoto, S., Shingu, A., Shirasu, Y. Teratogenicity and Dominant-Lethal Studies With O-Phenylphenol. J. Pesticide Sci. 3, 365-370, 1978
- 134. Kato, M. Reverse Mutation Assay of Preventol of Using Salmonella Typhimurium and Escherichia Coli. Research Lab., Jbc. Inc., Test No. 1097, 1989
- 135. Kawachi, T., Yahagi, T., Kada, T., Tazima, Y., Ishidate, M., Sasaki, M., Sugiyama, T. Cooperative Programme on Short-Term Assays For Carcinogenicity In Japan. Iarc Scientific Publ. 27, 323-330, 1980

- 136. Keller, B.J., Marsman, D.S., Popp, J.A., Thursman, R.G. Several Nongenotoxic Carcinogens Uncouple Mitochondrial Oxidative Phosphorilation. Biochim. Biophys. Acta 1102, 237-244, 1992
- 137. Kimmerle & Lorke; Study Report dated March 31, 1969
- 138. Kobayashi, H., Kabashima, J., Nakao, T. Effect of Sodium O-Phenylphenate (Opp-Na) On Alkaline Phosphatase in Rat Urine. Ann. Rep.Tokyo Metr. Res. Lab. P. H. 33, 467-469, 1982
- 139. Kojima, A., Fujita, H., Hiraga, K. Mutagenicity of O-Phenylphenol (OPP) in the Microbial System. Ann. Rep. Tokyo Metr. Res. Lab. P. H. 34, 319-324, 1983
- 140. Kojima, A., Hiraga, K. Mutagenicity of Citrus Fungicides In Microbial System. Ann. Rep. Tokyo Metr.Res. Lab. P.H. 29, 83-85, 1978
- 141. Kojima, K., Kanazawa, H.J.L., Ono, H. Long-Term Effects of Parathion, O-Phenylphenol and Penicillin-G Potassium on Immunological Properties and Intestinal Flora in BALB/C Mice The International Congress Of Toxicology Vii, July 2-6, 12, 1995 (Abstract)
- 142. Kolachana, P., Subrahmanyam, V.V., Eastmond, D.A., Smith, M.T. Metabolism of Phenylhydroquinone by Prostaglandin (H) Synthase: Possible Implications in Ophenylphenol Carcinogenesis. Carcinogenesis 12, 145-149, 1991
- 143. Kwok, E.S.C., Buchholz, B.A., Vogel, J.S., Turteltaub, K.W., Eastmond, D.A. Dose-Dependent Binding of Ortho-Phenylphenol to Protein but not DNA in the Urinary Bladder of Male F344 Rats. Toxicol. Appl. Pharmacol. 159, 18-24, 1999
- 144. Kwok, E.S.C., Eastmond, D.A. Effects of pH on Nonenzymatic Oxidation of Phenyl-Hydroquinone: Potential Role in Urinary Bladder Carcinogenesis Induced by O-Phenylphenol in Fischer 344 Rats. Chem. Res. Toxicol. 10, 742-749, 1997
- 145. La Via, M.F., La Via, D.S., Phenol Derivatives are Immunodepressive in Mice. Drug Chem. Toxicol. 2, 167-177, 1979a
- 146. La Via, M.F., Loose, L.D., La Via, D.S, Silberman, M.S. The Immunodepressive Effect of Phenol Derivatives. Adv. Exp. Med. Biol. 121, 523-538, 1979b
- 147. Lambert, A.C. Mechanisms of Genotoxicity Induced by the Ortho-Phenylphenol Metabolites Phenylhydroquinone and Phenyl-Benzoquinone. Master Thesis, Univ. Of California Riverside; 1992
- 148. Lambert, A.C., Eastmond, D.A. Genotoxic Effects of the O-Phenylphenol Metabolites Phenylhydroquinone and Phenylbenzoquinone In V79 Cells. Mutat. Res. 322, 243-256, 1994
- 149. Landry, T.D., Stebbins, K.E., Battjes, J.E. Ortho-Phenylphenol: Acute Aerosol Inhalation Toxicity Study in Fischer 344 Rats. Dow Chemical, Report No. K-001024-049, 1992

- 150. Loeser, E. Preventol O Extra: Untersuchungen zur Akuten Oralen Toxizität an Männlichen Wistar-Ratten. Bayer Ag, Report 1981
- 151. Luster, M.I., Dean, J.H., Boorman, G.A., Archer, D.L., Lauer, L., Lawson, L.D., Moore, J.A., Wilson, R.E. The Effects of Orthophenylphenol, Tris(2,3-Dichloropropyl)-Phosphate, and Cyclophosphamide on the Immune System and Host Susceptibility of Mice Following Subchronic Exposure. Toxicol. Appl. Pharmacol. 58, 252-261, 1981
- 152. Luster, M.I., Portier, C., Pait, D.G., Rosenthal, G.J., Germolec, D.R., Corsini, E., Blaylock, B.L., Pollock, P., Kouchi, Y., Craig, W., White, K.L., Munson, A., Comment, C.E. Risk Assessment in Immunotoxicology. Relationship Between Immune And Host Resistance Tests. Fund. Appl. Toxicol. 21, 71-82, 1993
- 153. Luster, M.I., Portier, C., Pait, D.G., White, K.L., Gennings C., Munson, A.E., Rosenthal, G.J. Risk Assessment in Immunotoxicology. I. Sensitivity and Predictability of Immune Tests. Fund. Appl. Toxicol. 18, 200-210, 1992
- 154. Macintosh, F.C. The Toxicity of Diphenyl and O-Phenylphenol. Analyst 70, 334-335, 1945
- 155. Maertins, T. Bayer Ag, Letter Report, 1988b
- 156. Maertins, T. Preventol ON: Untersuchungen zum Reiz-/Aetzpotential an Haut und Auge (Kaninchen) Nach OECD-Richtlinie No. 404 und 405. Bayer Ag, Report-No. 16951, 1988a
- 157. Marty et al., Ortho-Phenylphenol (OPP): Evaluation of Potential Estrogen, Androgen, or Thyroid Effects. Assessment Report; ID NS000106, Completion Date 13 November 2012 Lanxess Deutschland GmbH & DOW Benelux B.V.; 2-phenylphenol: Dossier according directive 91/414/EEC submission of confirmatory data; Dec. 2011
- 158. McMahon, R.E., Cline, J.C., Thompson, C.Z. Assay Of 855 Test Chemicals in Ten Tester Strains Using a New Modification of the Ames Test for Bacterial Mutagens. Cancer Res. 39, 682-693, 1979
- 159. McNett, D.A., Timchalk, C., Mendrala, A.L. Ortho-Phenylphenol: Metabolism of 14C-labelled OPP in B6C3F1 Mice snd Fischer 344 Rats. Dow Chemical, Report No. K-001024-060, 1997
- 160. Meiss, R., Heinrich, U., Himmels, S., Themann, H. A Morphometric Study of O-Phenylphenol Induced Alterations in the Rat Liver Following Oral or Subcutaneous Administration. Wissenschaft Und Umwelt, 4, 257-261, 1981
- 161. Mihail, F., Kimmerle, G. Preventol O und Preventol ON: Bestimmung der Inhalations-Toxizitaet. Bayer Ag, Report 1977

- 162. Mikuriya, H., Fujii, T., Hayashida, S., Kamiya, N., Hiraga, K. Examination of Dose-Related Enhancing Effects of Thiabendazole on Urinary Bladder Carcinogenesis by Sodium O-Phenylphenate in Rats. Ann. Rep. Tokyo. Metr. Res. Lab. P.H. 37, 407-410, 1986
- 163. Mikuriya, H., Fujii, T., Sasaki, M. Chronic Toxicity Test of O-Phenylphenol and Thiabendazole by Simultaneous Administration to Mice. Shokuhin Eiseigaku Zsasshi 33, 274-282, 1992
- 164. Mikuriya, H., Fujii, T., Yoneyama, M., Shimada, T., Sasaki, M. Toxicity of Thiabendazole and O-Phenylphenol by Simultaneous Administration to Rats for Six Weeks. Ann. Rep. Tokyo Metr. Res. Lab. P.H. 41, 245-254, 1990
- 165. Mikuriya, H., Fujii, T., Yoneyama, M., Yano, N., Yuzawa, K., Nagasawa, A., Sasaki, M. Toxicity of O-Phenylphenol by Dietary Administration to Mice For 52 Weeks. Ann. Rep. Tokyo. Metr. Res. Lab. P.H. 40, 281-288, 1989a
- 166. Mikuriya, H., Fujii, T., Yoneyama, M., Yano, N., Yuzawa, K., Nagasawa, A., Sasaki, M. Toxicity of Thiabendazole and O-Phenylphenol by Simultaneous Administration to Mice For 52 Weeks. Ann. Rep. Tokyo. Metr. Res. Lab. P.H. 40, 289-297, 1989b
- 167. Miller, D., Wheals, B.B., Beresford; N., Sumpter, J.P. Estrogenic Activity of Phenolic Additives Determined by an in vitro Yeast Bioassay. Environ. Health Perspect. 109, 133-138, 2001
- 168. Miyata, Y., Fukushima, S., Hirose, M., Masui, T., Ito, N. Short-Term Screening of Promoters of Bladder Carcinogenesis In N-Butyl-N-(4-Hydoxybutyl)Nitrosamine-Initiated; Unilaterally Ureter-Ligated Rats: Jpn. J. Cancer Res. 76, 828-834, 1985
- 169. Mori, S., Kurata, Y., Takeuchi, Y., Toyama, M., Makino, M., Fukushima, S. Influences of Strain and Diet on the Promoting Effects of Sodium L-Ascorbate in Two-Stage Urinary Bladder Carcinogenesis in Rats. Cancer Res. 47, 3492-3495, 1987
- 170. Morimoto, K., Fukuoka, M., Hasegawa, R., Tanaka, A., Takahashi, A., Hayashi, Y. DNA Damage in Urinary Bladder Epithelium of Male F344 Rats Treated With 2-Phenyl-1,4-Benzoquinone, One of the Non-Conjugated Urinary Metabolites of Sodium O-Phenylphenate. Jpn. J. Cancer Res. 78, 1027-1030, 1987
- 171. Morimoto, K., Sato, M., Fukuoka, M., Hasegawa, R., Takahashi, T., Tsuchiya, T., Tanaka, A., Takahashi, A., Hayashi, Y. Correlation Between the DNA Damage in Urinary Bladder Epithelium and the Urinary 2-Phenyl-1,4-Benzoquinone Levels from F344 Rats Fed Sodium O-Phenylphenate in the Diet. Carcinogenesis 10, 1823-1827, 1989

- 172. Moriya, M. Ohta, T., Watanabe, K., Miyazawa, T., Kato, K., Shirasu, Y. Further Mutagenicity Studies on Pesticides in Bacterial Reversion Assay Systems. Mutat. Res. 116, 185-216, 1983
- 173. Morpurgo, G., Bellincampi, D., Gualandi, G., Baldinelli, L., Crescenzi, O.S. Analysis of Mitotic Nondisjunction with Aspergillus Nidulans. Environ. Health Perspect. 31, 81-95, 1979
- 174. Mortelmans, K., Haworth, S., Lawlor, T., Speck, W., Tainer, B., Zeiger, E. Salmonella Mutagenicity Testing: Results from the Testing of 270 Chemicals. Environ. Mutagen. 8 (Suppl. 7), 1-119, 1986
- 175. Murata, M., Moriya, K., Inoue, S., Kawanishi, S. Oxidative Damage to Cellular and Isolated DNA by Metabolites of a Fungicide Orthophenylphenol. Carcinogenesis 20, 851-857, 1999
- 176. Nagai, F., Nakao, T. Changes in Enzyme Activities in the Urine and Tissues of Rats Fed Sodium O-Phenylphenate. Food Chem. Toxic. 22, 361-364, 1984
- 177. Nagai, F., Nakao, T. Changes of Enzyme Activities in Rat Urine and Tissues by Administration of Different Concentrations of O-Phenylphenol Sodium Salt (OPPNa) in Diet. Ann. Rep. Tokyo Metr. Res. Lab. P. H. 34, 305-308, 1983
- 178. Nagai, F., Nakao, T. Effect of O-Phenylphenol Sodium Salt (OPP-Na) on Rats Enzymes in vivo. Ann. Rep. Tokyo Metr. Res. Lab. P. H. 32, 57-60, 1981
- 179. Nagai, F., Ushiyama, K., Satoh, K., Kano, I. DNA Cleavage by Phenylhydroquinone: The Major Metabolite of a Fungicide O-Phenylphenol. Chem.-Biol. Interactions 76, 163-179, 1990
- 180. Nagai, F., Ushiyama, K., Satoh, K., Kasai, H., Kano, I. Formation of 8-Hydroxydeoxyguanosine in Calf Thymus DNA Treated in vitro with Phenylhydroquinone, the Major Metabolite of O-Phenylphenol. Carcinogenesis 16, 837-840, 1995
- 181. Nagai, F., Ushiyama. K., Satoh, K., Kano, I. Modification of DNA Base by OPP and its Metabolites. Ann. Rep. Tokyo Metr. Res. Lab. P.H. 39, 292-294, 1988
- 182. Nakagawa, A., Nakao, T., Nakao, M. Altered Levels of Cyclic Nucleotides in F344 Rats Fed Sodium O-Phenylphenate. Food Chem. Toxic. 22, 217-221, 1984
- 183. Nakagawa, Y., Moldéus, P., Moore, G.A. Cytotoxicity of Ortho-Phenylphenol in Isolated Rat Hepatocytes. Biochem. Pharmacol. 43, 159-165, 1992a
- 184. Nakagawa, Y., Moore, G.A. Cytotoxic Effects of Postharvest Fungicides, Ortho-Phenylphenol, Thiabendazole and Imazalil, on Isolated Rat Hepatocytes. Life Sciences 57, 1433-1440, 1995

- 185. Nakagawa, Y., Tayama, S. Effect of Buthionine Sulfoximine on Orthophenylphenol-Induced Hepato- And Nephrotoxic Potential in Male Rats. Arch. Toxicol. 62, 452-457, 1988
- 186. Nakagawa, Y., Tayama, S. Formation of Ortho-Phenylphenol Glutathione Conjugates in the Rat Liver. Xenobiotica 19, 499-507, 1989
- 187. Nakagawa, Y., Tayama, S. Induction of 8-Hydroxy-2´-Deoxyguanosine in Cho-K1 Cells Exposed to Phenyl-Hydroquinone, a Metabolite of Ortho-Phenylphenol. Cancer Lett. 101, 227-232, 1996
- 188. Nakagawa, Y., Tayama, S., Moldéus, P., Moore, G.A. Relationship Between Metabolism and Cytotoxicity of Ortho-Phenylphenol in Isolated Rat Hepatocytes. Biochem. Pharmacol. 43, 1431-1437, 1992b
- 189. Nakagawa, Y., Tayama, S., Moore, G., Moldéus, P. Cytotoxic Effects of Biphenyl and Hydroxybiphenyls on Isolated Rat Hepatocytes. Biochem. Pharmacol. 45, 1959-1965, 1993
- 190. Nakamura, K., Iguchi, S., Ikeada. T., Hiraga, K. Subacute Toxicity of O-Phenylphenol by Food Administration to Male Rats. Ann. Rep. Tokyo Metr. Lab. P. H. 32, 33-39, 1981
- 191. Nakamura, K., Ikeda, T., Iguchi, S., Hiraga, K. Toxicity of 2-Phenylphenol (OPP) by Dietary Administration to Male Rats For 91 Weeks. Ann. Rep. Tokyo Metr. Res. Lab. P. H. 33, 434-443, 1982
- 192. Nakao, T., Ushiyama, K., Kabashima, J., Nagai, F., Nakagawa, A., Ohno, T., Ichikawa, H., Kobayashi, H., Hiraga, K. The Metabolic Profile of Sodium O-Phenylphenate after Subchronic Oral Administration to Rats. Food Chem. Toxic. 21, 325-329, 1983
- 193. Narayan, S., Roy, D. Changes in Protein and Nonprotein Thiol Contents in Bladder, Kidney and Liver of Mice by the Pesticide Sodium-O-Phenylphenol and their Possible Role in Cellular Toxicity. Biochem. Int. 26, 191-198, 1992
- 194. Nawai, S., Yoshida, S., Nakao, T., Hiraga, K. Effect of O-Phenylphenol Incubated With S-9 Mix Induced by Phenobarbital or 3-Methylcholanthrene on Chromosomes in Cho-K1 Cells. Ann. Rep. Tokyo Metr. Res. Lab. P.H. 33, 480-483, 1982
- 195. Nawai, S., Yoshida, S., Nakao, T., Hiraga, K. Examination of Mutagens by Induced Sister Chromatid Exchange (SCE): Test of two Fungicides by Induced SCE, In Vitro. Ann. Rep. Tokyo Metr. Res. Lab. P.H. 30, 51-53, 1979
- 196. Neumann, H.-G., Bitsch, A., Kloehn P.-C. The Dual Role of 2-Acetylfluorene in Hepatocarcinogenesis: Specific Targets for Initiation and Promotion. Mut. Res. 376, 169-176, 1997

- 197. Nishioka, H., Ogasawara, H. Mutagenicity Testing for Diphenyl Derivatives in Bacterial Systems. Mutat. Res. 54, 248-249, 1979 (Abstract)
- 198. Norris, J. M. Eye Irritation Test Conducted on Dowicide 1 Preservative. Dow Chemical, Report 1971
- 199. NTIS = National Technical Information Service AS Ntis-Pb88-161989, 1988 Environmental Criteria and Assessment Office Health and Environmental Effects Profile For 2-Phenylphenol.
- 200. NTIS = National Technical Information Service Evaluation Of Carcinogenic, Teratogenic and Mutagenic Activities of Selected Pesticides and Industrial Chemicals. Vol. 1, Carcinogenic Study, Washington Dc, Us Department Of Commerce, 87, 1968
- 201. NTP National Toxicology Program Public Health Service, NTP-85-055, 1985
- 202. NTP Technical Report on the Toxicology and Carcinogenesis Studies of Ortho-Phenylphenol (CAS No. 90-43-7) alone and with 7,12-Dimethylbenz(A)Anthracene (CAS No. 57-97-6) in Swiss Cd-1 Mice Dermal Studies). NIH Publication No. 85-2557, NTP 84-099, US Department of Health and Human Services, NTP TR 301, 1986 (Also Cited As Luster, M., 1985 In Fao/Who Evaluations 1985)
- 203. OECD Environmental Health And Safety Publications Appraisal of Test Methods For Sex-Hormone Disrupting Chemicals. Series On Testing And Assessment, Paris, May 2001
- 204. Oehme, F.W. Comparative Toxicity of O-Phenylphenol and O-Phenylphenol-Containing Disinfectant. Toxicol. Appl. Pharmacol. 19, 412, 1971 (Abstract)
- 205. Oehme, F.W., Smith, T.H. The Metabolism and Urinary Excretion of O-Phenylphenol in Dogs and Cats. Toxicol. Appl. Pharmacol. 22, 292, 1972 (Abstract)
- 206. Ogata, A., Ando, H., Kubo, Y., Hiraga, K. Acute Oral Toxicity of Sodium O-Phenylphenate (Opp-Na) In Mice. Ann. Rep. Tokyo Metr. Res. Lab. P. H. 30, 54-56, 1979
- 207. Ogata, A., Ando, H., Kubo, Y., Hiraga, K. Dominant Lethal Tests of Long-Term Administration with Sodium O-Phenylphenol (Opp-Na) in Rats. Ann. Rep. Tokyo Metr. Res. Lab. P. H. 31, 17-19, 1980
- 208. Ogata, A., Ando, H., Kubo, Y., Hiraga, K. Teratological Tests of O-Phenylphenol (OPP) and Sodium O-Phenylphenol (Opp-Na) in Mice. Ann. Rep. Tokyo Metr. Res. Lab. P. H. 29, 89-96, 1978b
- 209. Ogata, A., Yoshida, S., Nawai, S., Ando, H., Kubo, Y., Hiraga, K., Masubuchi, M. Dominant Lethal Tests of Long-Term Administration with Sodium O-Phenylphenol (OPP-Na) in Mice. Ann. Rep. Tokyo Metr. Res. Lab. P. H. 29, 99-103, 1978a

- 210. Ohtsuki, K., Kabashima, J., Nakao, T. Urinary Metabolites of O-Phenylphenol Sodium Salt By Oral Administration. Ann. Rep. Tokyo Metr. Res. Lab. P. H. 32, 51-52, 1981
- 211. Okuda, M. Pathological Analysis of the Carcinogenic Effect of Sodium O-Phenylphenol and O-Phenylphenol in the Urinary Bladder of Rats and Mice. J. Nagoya City Univ. Med. Ass. 37, 157-184, 1986
- 212. Otoshi, T., Iwata, H., Yamamoto, S., Murai, T., Yamaguchi, S., Matsui-Yuasa, I., Otani, S., Fukushima, S. Severity of Promotion by Sodium Salts of Succinic Acid in Rat Urinary Bladder Carcinogenesis Correlates with Sodium Ion Concentration Under Conditions of Equal Urinary pH. Carcinogenesis 14, 2277-2281, 1993
- 213. Pagano, G., Cipollaro, M., Corsale, G., Della Morte, R., Esposito, A., Giordano, G.G., Micallo, G., Quinto, I., Staiano, N. Comparative Toxicity of Diphenyl, Diphenyl Ether, and Some of their Derivatives. Med. Biol. Environ. 16, 291-297, 1988
- 214. Pathak, D.N., Roy, D. Examination of Microsomal Cytochrome P450-Catalyzed in Vitro Activation of O-Phenylphenol to DNA Binding Metabolite(S) By 32p-Postlabeling Technique. Carcinogenesis 13, 1593-1597, 1992a
- 215. Pathak, D.N., Roy, D. In vivo Genotoxicity of Sodium Ortho-Phenylphenol: Phenylbenzoquinone is one of the DNA binding Metabolite(s) of Sodium Ortho-Phenylphenol. Mutat. Res. 286, 309-319, 1993
- 216. Pathak, D.N., Roy, D. Mechanism of Genotoxicity of O-Phenylphenol (OPP) in vivo. Proc. Am. Assoc. Cancer Res. 33, 141, 1992b (Abstract)
- 217. Pathak, D.N., Roy, D. Mechanisms of Genotoxicity of O-Phenylphenol in vitro: Covalent Modification to DNA by Phenyl-2,5´-P-Quinone, A Reactive Metabolite of O-Phenylphenol. Toxicologist 12, 252, 1992c (Abstract)
- 218. Pauluhn, J. Preventol on Extra: Untersuchungen auf Haut- und Schleimhautreizende Wirkung. Bayer Ag, Report 1983
- 219. Probst, G.S., Mcmahon, R.E., Hill, L.E., Thompson, C.Z., Epp, J.K., Neal, S.B. Chemically Induced Unscheduled DNA Synthesis in Primary Rat Hepatocyte Cultures: A Comparison With Bacterial Mutagenicity Using 218 Compounds. Environ. Mut. 3, 11-32, 1981
- 220. Quast, J.F., Mcguirk, R.J. Ortho-Phenylphenol:Two-Year Dietary Chronic Toxicity/Oncogenicity Study in B6C3F1 Mice. Dow Chemical Comp., Bayer-Miles, Report Code K-001024-047, 1995
- 221. Quast, J.F., Mcguirk, R.J., Kociba, R.J. Results of a Two-Year Dietary Toxicity/Oncogenicity Study of Ortho-Phenylphenol (OPP) In B6C3F1 Mice. Fundam. Appl. Toxicol. 36, 1997, 341 (Abstract)

- 222. Rachofsky, M.A., Oehme, F.W. Comparative And Age-Related Pharmocodynamics For Single And Multiple Doses of O-Phenylphenol. Toxicol. Appl. Pharmacol. 37, 93, 1976b (Abstract)
- 223. Rachofsky, M.A., Oehme, F.W. Distribution, Kinetics, and Tissue Residues of O-Phenylphenol in the Dog. Trans. Kansas Acad. Sci. 78, 23-24, 1976a (Abstract)
- 224. REACH Dossier 2-phenylphenol, 2010 ECHA website: http://echa.europa.eu
- 225. Reed, H.W.B. Alkylphenols. Kirk-Othmer Encyclopedia of Chemical Technology, 3rd Ed., Vol. 2, Grayson, M. And Eckroth, D., Eds., John Wiley and Sons, Inc., Ny. Pp 87-89, 1978
- 226. Rehmann, K., Schramm, K.-W., Kettrup, A.A. Applicability of a Yeast Oestrogen Screen for the Detection Of Oestrogen-Like Activities in Environmental Samples. Chemosphere 38, 3303-3312, 1999
- 227. Reitz, R.H., Fox, T.R., Quast, J.F., Hermann, E.A., Watanabe, P.G. Molecular Mechanisms Involved in the Toxicity of Orthophenylphenol and its Sodium Salt. Chem.-Biol. Interactions 43, 99-119, 1983
- 228. Reitz, R.H., Fox, T.R., Quast, J.F., Hermann, E.A., Watanabe, P.G. Biochemical Factors Involved in the Effects of Orthophenylphenol (OPP) and Sodium Orthophenylphenate (SOPP) on the Urinary Tract of Male F344 Rats. Toxicol. Appl. Pharmacol. 73, 345-349, 1984
- 229. Robenek, H., Meiss, R., Themann, H., Himmels, S. A Correlated Thin Section And Freeze-Fracture Study Of O-Phenylphenol-Induced Alterations in the Rat Liver. Expl. Cell Biol. 48, 404-420, 1980
- 230. Rodent Bladder Carcinogenesis Group. Urinary Bladder Carcinogenesis: Implications for Risk Assessment. Food Chem. Toxicol. 33, 797-802, 1995
- 231. Romanoski, C.A., Eastmond, D.A. Arachidonic Acid-Dependent Induction of Micronuclei in V79 Cells by Phenylhydroquinone, A Metabolite of the Fungicide O-Phenylphenol. Toxicologist 12, 56, 1992 (Abstract)
- 232. Routledge, E.J., Sumpter, J.P. Structural Features of Alkylphenolic Chemicals Associated With Estrogenic Activity. J. Biol. Chem. 272, 3280-3288, 1997
- 233. Roy, D. Cytochrome P-450 Catalyzed Redox Cycling of Orthophenylphenol. Biochem. Int. 22, 849-857, 1990
- 234. Sakai, A., Miyata, N., Takahashi, A. Initiating Activity of Quinones in the Two-Stage Transformation of BALB/3T3 Cells. Carcinogenesis 16, 477-481, 1995
- 235. Salminen, E., Salminen, S. Urinary Excretion of Orally Administered Oxalic Acid in Saccharin and O-Phenylphenol-Fed Nmri Mice. Urol. Int. 41, 88-90, 1986

- 236. San, H.C., Springfield, K.A. O-Phenylphenol Salmonella/Mammalian-Microsome Plate Incorporation Mutagenicity Assay (Ames Test). Microbiological Associates Inc., (Lab. Study Number C141.501017); 1989a
- 237. San, H.C., Springfield, K.A. O-Phenylphenol, Sodium Salt Tetrahydrate Salmonella/Mammalian-Microsome Plate Incorporation Mutagenicity Assay (Ames test). Microbiological Associates Inc., (Lab. Study Number C142.501017); 1989b
- 238. Sasaki, M., Nakao, T. The Effects of O-Phenylphenol On The Immune Response In Vitro. Ann. Rep. Tokyo Metro. Res. Lab. P.H. 29, 109-111, 1978
- 239. Sasaki, M., Ueno, S., Miyamae, Y., Ohta, T., Tsuda, S. Detection of Organ Specific Genotoxicity by Alkaline Single Cell Gel Electrophoresis Assay With Mouse Multiple Organs. Environ. Mutagen. Res. 20, 51-62, 1998
- 240. Sasaki, Y. The Comet Assay With 8 Mouse Organs: Results With 39 Currently Used Food Additives, Mutation Research 519, 103-119, 2002
- 241. Sasaki, Y.F. Saga, A., Akasaka, M., Yoshida, K., Nishidate, E., Su, Y.Q., Matsusaka, N., Tsuda, S. In vivo Genotoxicity of Ortho-Phenylphenol, Biphenyl, And Thiabendazole Detected in Multiple Mouse Organs by the Alkaline Single Cell Gel Electrophoresis Assay. Mutat. Res. 395, 189-198, 1997
- 242. Sato, H., Toyoda, K., Takamura, N., Furukawa, F., Hasegawa, R., Fukuoka, M., Imaida, K., Takahashi, M., Hayashi, Y. Effects of 2-Phenyl-1,4-Benzoquinone and 2,5-Dihydroxybiphenyl on Two-Stage Mouse Skin Carcinogenesis. Cancer Lett. 55, 233-238, 1990
- 243. Sato, M., Tanaka, A., Tsuchiya, T., Yamaha, T., Nakaura, S., Tanaka, T. Excretion, Distribution and Metabolic Fate of Sodium O-Phenylphenate and O-Phenylphenol in the Rat. J. Food Hyg. Soc. Japan 29, 7-12, 1988
- 244. Savides, M.C., Oehme, F.W. Urinary Metabolism Of Orally Administered Ortho-Phenyl-Phenol in Dogs and Cats. Toxicology 17, 355-363, 1980
- 245. Schewe, T., Markgraf, K., Schewe, C., Fischer, S., Getter, R., Mayer, M. Stoffwechselschädigung Menschlicher Hautzellen Durch Orthophenylphenol (OPP). Melliand Textilberichte 9, 631-632, 1997
- 246. Schnuch, A., Geier, J., Uter, W., Frosch, P.J. Patch Testing With Preservatives, Antimicrobials and Industrial Biocides. Results from a Multicentre Study. Brit. J. Dermatol. 138, 467-476, 1998
- 247. Schreiber, G. Prüfung Von Preventol O Extra auf Primäre Hautreizwirkung. Fraunhofer-Institut für Toxikologie und Aerosolforschung, 1981a
- 248. Schreiber, G. Prüfung von Preventol O Extra auf Schleimhautreizwirkung. Fraunhofer-Institut für Toxikologie Und Aerosolforschung, 1981b

- 249. Schumann, R. Tagung Der Kommission für Kosmetische Mittel Des BGVV; Bericht über die 59. Sitzung am 7. Dezember in Berlin. Bundesgesundheitsbl. Gesundheitsforsch. Gesundheits-Schutz 43, 386-387, 2000
- 250. Sekihashi K., Comperative Investigations of Multiple Organs of Mice and Rats in the Comet Assay Mutat. Res, 517, 53-74, 2002
- 251. Selim, S. A Single Dose Open Label Study to Investigate the Absorption and Excretion of 14c/13c-Labeled Orthophenylphenol Formulation After Dermal Application to Healthy Volunteers. Pharma Bio-Research Clinics Bv, Assen, Netherlands, Report No. P0995002, 1996
- 252. Shibata, A., Hagiwara, A., Fukushima, S., Ito, N. Thirteen Weeks Subacute Oral Study with Sodium Ortho-Phenylphenate (OPP-Na) In B6C3F1 Mice. J. Toxic. Sci. 6, 257, 1981 (Abstract)
- 253. Shibata, M., Kagawa, M., Kawabe, M., Hagiwara, A., Fukushima, S. Comparative Promoting Activities of Phosphate Salts on Rat Two-Stage Bladder Carcinogenesis Under Conditions Of Equivalent Urinary Na+ or K+ Levels. Teratogenesis, Carcinogenesis, And Mutagenesis 11, 305-316, 1991
- 254. Shibata, M., Tamano, S., Shirai, T., Kawabe, M., Fukushima, S. Inorganic Alkalizers and Acidifiers Under Conditions of High Urinary Na+ Or K+ on Cell Proliferation and Two-Stage Carcinogenesis in the Rat Bladder. Jpn. J. Cancer Res. 83, 821-829, 1992
- 255. Shibata, M.-A., Hagiwara, A., Tamano, S., Fukushima, S., Ito, N. Subchronic Toxicity Study of Sodium O-Phenylphenate in Mice. Toxicol. Letters 25, 239-246, 1985
- 256. Shibata, M.-A., Nakanishi, K., Shibata, M., Masui, T., Miyata, Y., Ito, N. Promoting Effect of Sodium Chloride in 2-Stage Urinary Bladder Carcinogenesis in Rats Initiated By N-Butyl-N-(4-Hydroxybutyl)-Nitrosamine. Urol. Res. 14, 201-206, 1986
- 257. Shibata, M.-A., Tamano, S., Kurata, Y., Hagiwara, A., Fukushima, S. Participation of Urinary Na+, K+, pH and L-Ascorbic Acid in the Proliferative Response of the Bladder Epithelium After the Oral Administration of Various Salts and/or Ascorbic Acid To Rats. Food Chem. Toxicol. 27, 403-413, 1989a
- 258. Shibata, M.-A., Tanaka, H., Yamada, M., Tamano, S., Fukushima, S. Proliferative Response of Renal Pelvic Epithelium in Rats to Oral Administration of Orthophenylphenol, Sodium Ortho-Phenylphenate and Diphenyl. Cancer Lett. 48, 19-28, 1989b
- 259. Shibata, M.-A., Yamada, M., Tanaka, H., Kagawa, M., Fukushima, S. Changes in Urine Composition, Bladder Epithelial Morphology, and DNA Synthesis in Male F344

- Rats in Response to Ingestion of Bladder Tumor Promotors. Toxicol. Appl. Pharmacol. 99, 37-49, 1989c
- 260. Shioya, S., Nagami-Oguihara, R., Oguihara, S., Kimura, T., Imaida, K. Roles of Bladder Distension, Urinary Ph And Urinary Sodium Ion Concentration in Cell Proliferation of Urinary Bladder Epithelium In Rats Ingesting Sodium Salts. Food Chem. Toxicol. 32, 165-171, 1994
- 261. Shirasu, Y., Moriya, M., Kato, K., Tezuka, H., Henmi, R., Shingu, A., Kaneda, M., Teramoto, S. Mutagenicity Testing on O-Phenylphenol. Mutat. Res. 54, 227, 1978 (Abstract)
- 262. Shirasu, Y., Moriya, M., Tezuka, H., Teramoto, S., Ohta, T., Inoue, T. Mutagenicity Screening Studies On Pesticides. International Conference On Environmental Mutagens 3, 331-335, 1982
- 263. Smith, R.A., Christenson, W.R., Bartels, M.J., Arnold, L.L., St. John, M.K., Cano, M., Garland, E.M., Lake, S.G., Wahle, B.S., Mcnett, D.A., Cohen, S.M. Urinary Physiologic and Chemical Metabolic Effects on the Urothelial Cytotoxicity and Potential DNA Adducts of O-Phenylphenol in Male Rats. Toxicol. Appl. Pharmacol. 150, 402-413, 1998
- 264. Soto, A.M., Fernandez M.F., Luizzi M.F., Karasko, A.S.O., Sonnenschein, C. Developing a Marker of Exposure to Xenoestrogen Mixtures in Human Serum. Environ Health Perspect 105 (Suppl 3), 647-654, 1997
- 265. St.John, M.K., Arnold, L.L., Anderson, T., Cano, M., Johansson, S.L., Cohen, S.M. Dietary Effects of Ortho-Phenylphenol and Sodium Ortho-Phenylphenate on Rat Urothelium. Toxicol. Sci. 59, 346-351, 2001
- 266. Storrs, F.J., Rosenthal, L.E., Adams, R.M., Clendenning, W., Emmett, E.A., Fisher, A.A., Larsen, W.G., Maibach, H.I., Rietschel, R.L., Schorr, W.F., Taylor, J.S. Prevalence And Relevance Of Allergic Reactions in Patients Patch Tested in North America 1984 To 1985. J. Am. Acad. Dermatol. 20, 1038-1045, 1989
- 267. Stouten, H. Toxicological Profile for O-Phenylphenol and its Sodium Salt. J. Appl. Toxicol. 18, 261-270, 1998
- 268. Suberg, H. Preventol O Extra (OPP): Untersuchung auf Primäre Reiz-/Aetzwirkung an der Kaninchenhaut. Bayer Ag, Inst. F. Toxikologie, Report 1983
- 269. Sugihara, N., Shimomichi, K., Furuno, K. Cytotoxicity of Food Preservatives in Cultured Rat Hepatocytes Loaded With Linolenic Acid. Toxicology 120, 29-36, 1997
- 270. Suzuki, H., Nakao, T., Hiraga, K. Mutagenicity of Orthophenylphenol (OPP) on a Human Clonal Cell Line. Ann. Rep. Tokyo Metr. Res. Lab. P.H. 35, 399-400, 1984
- 271. Suzuki, H., Suzuki, N., Sasaki, M., Hiraga, K. Orthophenylphenol Mutagenicity in A Human Cell Strain. Mutat. Res. 156, 123-127, 1985

- 272. Tada, Y., Fujitani, T., Yano, N.Y., Yuzawa, K., Nagasawa, A., Aoki, N., Ogata, A., Yoneyama, M. Chronic Toxicity Of Thiabendazole (TBZ( in CD-1) Mice. Toxicology 169, 163-176, 2001
- 273. Tadi-Uppala. P., Hasegawa, L., Rupa, D.S., Eastmond, D.A. Detection of Micronuclei and Cell Proliferation in the Rat Bladder Induced by the Fungicides Ophenylphenol and Sodium Ortho-Phenylphenate. Carcinogenesis 37, 127-128, 1996 (Abstract)
- 274. Takahashi, T., Sato, H., Toyoda, K., Furukawa, F., Imaida, K., Hasegawa, R., Hayashi, Y. Sodium O-Phenlyphenate (OPP-Na) Promotes Skin Carcinogenesis in CD-1 Female Mice Initiated with 7,12-Dimethylbenz[A]-Anthracene. Carcinogenesis 10, 1163-1167, 1989
- 275. Taniguchi, Y., Morimoto, J., Okada, K., Imai, S., Tsubura, Y. Toxicological Study of O-Phenylphenol (OPP) in Mice: I. Acute Oral Toxicity in DDY Mouse. J. Nara Med. Ass. 32, 425-429, 1981a
- 276. Taniguchi, Y., Morimoto, J., Okada, K., Imai, S., Tsubura, Y. Toxicological Study of Sodium Orthophenylphenate (Opp-Na) in Rats. Acute Oral Toxicity in Fischer 344 DUCRJJ Rats. J. Nara. Med. Ass. 32, 709-714, 1981b
- 277. Tayama, K., Hiraga, K. Depigmentation Caused by a Single Oral Administration of O-Phenylphenol (OPP) in Ordinary and Textile Hair Of C57BL/6N Mice. Ann. Rep. Tokyo Metr. Res. Lab. P.H. 35, 401-406, 1984
- 278. Tayama, K., Iguchi, S., Hiraga, K. Acute Oral Toxicity of O-Phenylphenol in Rats. Ann. Rep. Tokyo Metr. Res. Lab. P.H. 31, 1-6, 1980
- 279. Tayama, K., Iguchi, S., Hiraga, K. Acute Oral Toxicity of Sodium O-Phenylphenol (OPP-Na) in Rats. Ann. Rep. Tokyo Metr. Res. Lab. P.H. 30, 57-65, 1979
- 280. Tayama, K., Iguchi, S., Sasaki, M., Hiraga, K. Acute Oral Toxicity of O-Phenylphenol (Opp) in Mice. Ann. Rep. Tokyo Metr. Res. Lab. P.H. 34, 325-328, 1983a
- 281. Tayama, K., Nakamura, K., Kamiya, N., Hiraga, K. Urinalyses of Male F344/DUCRJ Rats Fed with Sodium O-Phenylphenate (OPP-Na). Ann. Rep. Tokyo Metr. Res. Lab. P. H. 35, 425-430, 1984
- 282. Tayama, S., Ichikawa, H. Effects of Phenylphenols on Chromosomes in CHO-K1 Cells. Ann. Rep. Tokyo Metr. Res. Lab. P. H. 38, 388-391, 1987
- 283. Tayama, S., Kamiya, N., Nakagawa, Y. Genotoxic Effects of O-Phenylphenol Metabolites in CHO-K1 Cells. Mutat. Res. 223, 23-33, 1989
- 284. Tayama, S., Kamiya, N., Nakao, T., Hiraga, K. Detection of O-Phenylphenol (OPP) and the Activated Metabolites with S-9 mix by HPLC and the Effect of These

- on Induction of SCE in CHO-K1 Cells. Ann. Rep. Tokyo Metr. Res. Lab. P.H. 34, 312-314, 1983b
- 285. Tayama, S., Nakagawa, Y. Effect of Scavangers of Active Oxygen Species on Cell Damage Caused in CHO-K1 Cells by Phenylhydroquinone, an O-Phenylphenol Metabolite. Mutat. Res. 324, 121-131, 1994
- 286. Tayama, S., Nakagawa, Y. Sulfhydryl Compounds Inhibit the Cyto- and Genotoxicity of O-Phenylphenol Metabolites in CHOK1 Cells. Mutat. Res. 259, 1-12, 1991
- 287. Tayama-Nawai, S., Yoshida, S., Nakao, T., Hiraga, K. Induction of Chromosome Aberrations and Sister-Chromatid Exchanges in CHO-K1 Cells by O-Phenylphenol. Mutat. Res. 141, 95-99, 1984
- 288. Thalacker, F.W. Nature of the Residue of 14C-Orthophenylphenol in Lactating Goats. Covance Laboratories Inc., Laboratory Project Identification Chw 6578-105, 1997
- 289. Thyssen, J. Preventol O Extra: Gewerbetoxikologische Untersuchungen. Bayer Ag, Report No. 10541, 1982
- 290. Timchalk, C., Selim, S., Sangha, G., Bartels, M.J. The Pharmacokinetics AMD Metabolism of 14C/13C-Labeled Ortho-Phenylphenol Formation Following Dermal Application to Human Volunteers. Hum. Exp. Toxicol. 17, 411-417, 1998
- 291. Timchalk, K. 14c-Orthophenylphenol: Pharmacokinetics Following Dermal Application in Male Human Volunteers. Dow Chemical, Report No. K-001024-064, 1996 page 134 of 146
- 292. Trotz, S.I., Pitts, J.J. Industrial Antimicrobial Agents. Kirk-Othmer Encyclopedia of Chemical Technology, 3rd Ed., Vol. 13, Grayson M., Eckroth, D., Eds., John Wiley and Sons, Inc., Ny. Pp 226-229, 1981
- 293. Tuer, W.F., James, W.D., Summers, R.J. Contact Urticaria to O-Phenylphenate. Annals Allergy 56, 19-21, 1986
- 294. US EPA: Reregistration Eligibility Decision for 2-phenylphenol and Salts (Orthophenylphenol or OPP), July 28, 2006
- 295. Ushiyama, K., Kabashima, J., Nakao, T. Metabolism of 2-Phenylphenol (OPP) in Rats: Metabolic Profile of OPP in Rats Fed Dietary for Long Period. Ann. Rep. Tokyo Metr. Res. Lab. P.H. 33, 455-457, 1982
- 296. Ushiyama, K., Kabashima, J., Nakao, T. Metabolism of O-Phenylphenol Sodium Salt (OPP-Na) in Rats: Dose-Response of Metabolic Profile of OPP. Ann. Rep. Tokyo Metr. Res. Lab. P.H. 34, 297-298, 1983
- 297. Ushiyama, K., Nagai, F., Nakagawa, A., Kano, I. DNA Adduct Formation by O-Phenyphenol Metabolite in vivo and in vitro. Carcinogenesis 13, 1469-1473, 1992

- 298. Ushiyama, K., Sato, K., Nakagawa, A., Nagai, F., Kabashima, J., Kano, I., Nakao, T. The Metabolism of O-Phenylphenol Sodium Salt. Toxicol. Lett. 31 (Suppl.), 168, 1986 (Abstract)
- 299. Uter, W., Schaller, S., Bahmer, F.A., Brasch, J., Diepgen, T.L., Enders, F., Frosch, P.J., Fuchs, T., Henseler, T., Mueller, S., Peters, K.P., Przybilla, B., Schaller, J., Schnuch, A., Schulze-Dirks, A., Stary, A. Contact Allergy in Metal Workers A One-Year Analysis Based on Data Collected by the "Information Network of Dermatological Clinics" (IVDK) in Germany. Dermatosen, 41, 320-327, 1993
- 300. Uwagawa, S., Imaida, K., Tsuda, H., Masui, T., Ito, N. Marked Enhancing Potential of Prior N-Methyl-N-Nitrosourea (MNU) Treatment on Rat Tumorigenesis in Various Organs Induced by 6 Different Carcinogens. Toxicologist 8, 165, 1988 (Abstract)
- 301. Uwagawa, S., Tsuda, H., Inoue, T., Tagawa, Y., Aoki, T., Kagawa, M., Ogiso, T., Ito, N. Enhancing Potential of 6 Different Carcinogens on Multi-Organ Tumorigenesis After Initial Treatment with N-Methyl-N-Nitrosourea in Rats. Jpn. J. Cancer Res. 82, 1397-1405, 1991
- 302. Van Hecke, E. Contact Dermatitis to O-Phenylphenol in a Coolant. Contact Dermatitis 15, 46, 1986
- 303. Wahle, B.S., Christenson, W.R. Technical Grade Ortho-Phenylphenol: A Combined Chronic Toxicity/Oncogenicity Testing Study in the Rat. Bayer Corp., Report No. 92-272-Sc, 1996
- 304. Wahle, B.S., Christenson, W.R., Lake, S.G., Elcock, L.E., Moore, K.D., Sangha, G.K., Thyssen, J.H. Technical Grade Ortho-Phenylphenol: A Combined Chronic Toxicity/Oncogenicity Testing Study in the Rat. Fundam. Appl. Toxicol. 36, 1997 (Abstract)
- 305. Weckmann: Test Report: Dermatological skin-tolerance studies with Preventol O extra. Lanxess Deutschland GmbH Leverkusen, February 15, 2012
- 306. WHO/IPCS/00.4: Pesticide residues in food 1999. Toxicological evaluations: 2-Phenylphenol and its sodium salt, 201-237, 2000
- 307. WHO; World Health Organization. Pesticide Residues In Food 1999 WHO/PSC/00.4, 201-237, 2000
- 308. Wiebkin, P., Fry, J.R., Jones, C.A., Lowing, R.K., Bridges, J.W. The Metabolism of Biphenyl by Isolated Viable Rat Hepatocytes. Xenobiotica 6, 725-743, 1976
- 309. Wiebkin, P., Fry, J.R., Jones, C.A., Lowing, R.K., Bridges, J.W. Biphenyl Metabolism in Isolated Rat Hepatocytes: Effect of Induction and Nature of the Conjugates. Biochem. Pharmacol. 27, 1899-1907, 1978

- 310. Woodruff, R.C., Mason, J.M., Valencia, R., Zimmering, S. Chemical Mutagenesis Testing in Drosophila. V. Results of 53 Coded Compounds Tested for the National Toxicology Program. Environ. Mut. 7, 677-702, 1985
- 311. WRC: EUROPEAN COMMISSION STUDY ON THE SCIENTIFIC EVALUATION OF 12 SUBSTANCES IN THE CONTEXT OF ENDOCRINE DISRUPTER PRIORITY LIST OF ACTIONS. Authors: I Johnson and P Harvey; WRc-NSF Ref: UC 6052, NOVEMBER 2002
- 312. Yamazaki, H., Yamaguchi, T., Yamauchi, A., Kakiuchi, Y. Food Additives on Acceptable Daily Intake (ADI) Level Affect the Agonist Induced Platelet Activation.

  I. Antioxidants And Preservatives: Chemosphere 29, 1293-1299, 1994
- 313. Yanagisawa, F., Ueda, Y., Tsuchida, M. Acute Intoxication by O-Phenylphenol. Jpn. J. Rural Med. 27, 632-633, 1978
- 314. Yoshida, S., Hiraga, K. Cytogenetic Studies on Rats Fed with O-Phenylphenol (OPP) and Sodium O-Phenylphenate (OPP-Na) for Long-Term. Ann. Rep. Tokyo Metr. Res. Lab. P.H. 33, 489-491, 1982
- 315. Yoshida, S., Nawai, S., Hiraga, K. Cytogenetic Studies of Sodium O-Phenylphenate (OPP-Na). Ann. Rep. Tokyo Metr. Res. Lab. P.H. 30, 44-47, 1979
- 316. Yoshida, S., Sasaki, M., Nakao, T., Hiraga, K. Chromosome Preparations from CHO-K1 Cells Used NUNC Plate. Ann. Rep. Tokyo Metr. Res. Lab. P.H. 32, 111-114, 1981
- 317. Zablotny, C.L., Breslin, W.J., Kociba, R.J. Developmental Toxicity of Ortho-Phenylphenol (OPP) in New Zealand White Rabbits. Toxicologist 12, 103, 1992 (Abstract)
- 318. Zablotny, C.L., Breslin, W.J., Kociba, R.J. Ortho-Phenylphenol (Opp): Gavage Teratology Probe Study in New Zealand White Rabbits. Dow Chemical Report No. K-001024-044, 1991a
- 319. Zablotny, C.L., Breslin, W.J., Kociba, R.J. Ortho-Phenylphenol (Opp): Gavage Teratology Study in New Zealand White Rabbits. Dow Chemical Report No. K-001024-045, 1991b
- 320. Zempel, J.A., Szabo, J.R. Ortho-Phenylphenol: 21-Day Repeated Dermal Dose Study of Systemic Toxicity in Fischer 344 Rats. Dow Chemical Report No. K-001024-056, 1993

#### **SCCS References:**

Anses (2014):

https://www.anses.fr/en/content/anses-publishes-its-recommendations-reduce-exposure-five-substances-which-are-reproductive

https://www.anses.fr/en/documents/CHIM2009sa0331Ra-01.pdf (évaluation des risques sanitaires des substances reprotoxiques et/ou perturbatrices endocriniennes dans les produits de consommation : le o-phénylphénol (OPP) (n° CAS 90-43-7)).

https://www.anses.fr/en/documents/CHIM2009sa0331Ra-01-An01.pdf (Filières, usages et expositions liées à la présence de substances reprotoxiques et/ou perturbatrices endocriniennes dans les produits de consommation : le o-phénylphénol (OPP) (n° CAS 90-43-7)).

https://www.anses.fr/en/documents/CHIM2009sa0331Ra-01-An02.pdf (Profil toxicologique de l'o-phénylphénol (OPP) (n° CAS 90-43-7)).

Ghisari, M. and Bonefeld-Jorgensen, E.C. (2009): Effects of plasticizers and their mixtures on estrogen receptor and thyroid hormone functions. Toxicology Letters 189, 67-77.

Vinggaard, A.M., Körner, W., Lund, K.H., Bolz, U. And Petersen, J.H. (2000): Identification and Quantification of Estrogenic Compounds in Recycled and Virgin Paper for Household Use as Determined by an in Vitro Yeast Estrogen Screen and Chemical analysis. Chem. Res. Toxicol. 13, 1214 – 1222.

Kwok, E.S.C. and Silva, M. (2013): Re-evaluation of Developmental and Reproductive Toxicity of Ortho-Phenylphenol (OPP) and Sodium Ortho-Phenylphenate (SOPP). Cell and Developmental Biology 2:3. http://dx.doi.org/10.4172/2168-9296.1000123

EFSA (2008): Conclusion on Pesticide Peer Review. Peer review of the pesticide risk assessment of the active substance 2-phanylphenol (Question No EFSA-Q-2008-392). EFSA Scientific Report 217, 1-67.

Available online: <a href="http://www.efsa.europa.eu/de/efsajournal/doc/217r.pdf">http://www.efsa.europa.eu/de/efsajournal/doc/217r.pdf</a>.

US EPA (2006): Reregistration Eligibility Decision for 2-phenylphenol and Salts (Orthophenylphenol or OPP). Office of Prevention, Pesticide and Toxic Substances (7510C) 739-R-06-004).

Available online:

http://www.epa.gov/oppsrrd1/REDs/phenylphenol\_red.pdf. (corresponds to Ref. 294).

Cal EPA (2007): Ortho-Phenylphenol (OPP) and Sodium Ortho-Phenylphenate (SOPP). Risk Characterization Document Dietary Exposure.

Available online: <a href="http://www.cdpr.ca.gov/docs/risk/rcd/opp.pdf">http://www.cdpr.ca.gov/docs/risk/rcd/opp.pdf</a>.

European Commission (2002): Study on the scientific evaluation of 12 substances in the context of endocrine disrupter priority list of actions. (I. Johnson and P. Harvey, Authors). Report No.: UC 6052.

Petit, F., Le Goff, P., Cravédi, J.P., Valotaire, Y., and Pakdel, F. (1997): Two complementary bioassays for screening the estrogenic potency of xenobiotics: recombinant yeast for trout estrogen receptor and trout hepatocyte cultures. J. Mol. Endoc. 19, 321 – 335.

Imai, K., Yoshimura, S., Hashimoto, K., and Boorman, G.A. (1991): Effect of dietary restriction on age-associated pathological changes in Fischer 344 rats. In: ILIS Monographs: Biological effects of dietary restriction (Ed: Fishbein, L) p.87-98.

Niho, N., Shibutani, M., Toyoda, K., Sato, H., Hirose, A., Imaida, K., Takahashi, M., Hayashi, Y. and Hirose, M. (2002): Dose- and time-response studies of sodium ophenylphenate urinary bladder carcinogenicity in rats. Food Chem. Toxicol. 40, 715 – 722. Habicht, J. and Brune, K. (1983): Inhibition of prostaglandin E2 release by salicylates, benzoates and phenols: a quantitative structure-activity study. Journal of Pharmacy and Pharmacology 35, 718 – 723.

NTP (2007) Historical Controls Report All Routes And Vehicles. MICE, October 2007. http://ntp.niehs.nih.gov/files/HistCont-2007-10-17-Mice All Routes.pdf

Haseman J.K, Hailey J.R. and Morris, R.W. (1998): Spontaneous neoplasm incidences in Fischer 344 rats and B6C3F-1 mice in two-year carcinogenicity studies: A national toxicology program update. Toxicol Pathol. 26, 428-441.

Battershill, J.M. and Fielder, R.J. (1998): Mouse-specific carcinogens: an assessment of hazard and significance for validation of short-term carcinogenicity bioassays in transgenic mice. Hum. Exp. Toxicol. 17, 193-205.

# References obtained after public consultation:

US EPA (2015): EDSP: Weight of Evidence analysis of Potential Interaction with the Estrogen, Androgen or Thyroid Pathways; Chemical: o-Phenylphenol (o-PP) http://www2.epa.gov/sites/production/files/2015-06/documents/o-phenylphenol-064103\_2015-06-29\_txr0057146.pdf

# **References obtained in December 2017:**

Opinion of the German Federal Institute for Risk Assessment

## 6. GLOSSARY OF TERMS

See SCCS/1564/15, 9th Revision of the SCCS Notes of Guidance for the Testing of Cosmetic Ingredients and their Safety Evaluation – from page 144

## 7. LIST OF ABBREVIATIONS

See SCCS/1564/15, 9th Revision of the SCCS Notes of Guidance for the Testing of Cosmetic Ingredients and their Safety Evaluation – from page 144